

Sums of powers of binomials, their Apéry limits, and Franel's suspicions

Southern Regional Number Theory Conference
LSU

Armin Straub


Mar 13, 2022

University of South Alabama

Special thanks to NSF and NSA for supporting this conference.

CONJ $\pi, \zeta(3), \zeta(5), \dots$ are algebraically independent over \mathbb{Q} .

- Apéry (1978): $\zeta(3)$ is irrational
- Open: $\zeta(5)$ is irrational
- Open: $\zeta(3)$ is transcendental
- Open: $\zeta(3)/\pi^3$ is irrational



based on joint work(s) with:



Marc Chamberland
(Grinnell College)




Wadim Zudilin
(Radboud University)

The last shall be first: conclusions

CONJ
Frenel,
1895

The minimal recurrence for $A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$ has order $\lfloor \frac{s+1}{2} \rfloor$.



THM
Stoll '97


$A^{(s)}(n)$ satisfies a recurrence of order $\lfloor \frac{s+1}{2} \rfloor$.



The last shall be first: conclusions

CONJ
Frenel,
1895

The minimal recurrence for $A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$ has order $\lfloor \frac{s+1}{2} \rfloor$.



THM
Stoll '97

$A^{(s)}(n)$ satisfies a recurrence of order $\lfloor \frac{s+1}{2} \rfloor$.



$$\sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$


THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

The last shall be first: conclusions

CONJ
Franel,
1895

The minimal recurrence for $A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$ has order $\lfloor \frac{s+1}{2} \rfloor$.



THM
Stoll '97

$A^{(s)}(n)$ satisfies a recurrence of order $\lfloor \frac{s+1}{2} \rfloor$.



$$\sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover, $A_j^{(s)}(n)$ with $0 \leq 2j < s$ are linearly independent, so that any telescoping recurrence has order at least $\lfloor \frac{s+1}{2} \rfloor$.

A victory for the French peasant...*

- The **Apéry numbers**

1, 5, 73, 1445, ...

satisfy

$$A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$$

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

THM
Apéry '78

$\zeta(3) = \sum_{n=1}^{\infty} \frac{1}{n^3}$ is irrational.



* Someone's "sour comment" after Henri Cohen's report on Apéry's proof at the '78 ICM in Helsinki.

A victory for the French peasant...*

- The **Apéry numbers**

1, 5, 73, 1445, ...

satisfy

$$A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$$

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$



THM
Apéry '78

$\zeta(3) = \sum_{n=1}^{\infty} \frac{1}{n^3}$ is irrational.

proof The same recurrence is satisfied by the “near”-integers

$$B(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left(\sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right).$$

Then, $\frac{B(n)}{A(n)} \rightarrow \zeta(3)$. But too fast for $\zeta(3)$ to be rational. \square

* Someone's “sour comment” after Henri Cohen's report on Apéry's proof at the '78 ICM in Helsinki.

A victory for the French peasant...*

- The **Apéry numbers**

1, 5, 73, 1445, ...

satisfy

$$A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$$

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

THM
Apéry '78

$\zeta(3) = \sum_{n=1}^{\infty} \frac{1}{n^3}$ is irrational.



“After a few days of fruitless effort the specific problem was mentioned to Don Zagier (Bonn), and with *irritating speed* he showed that indeed the sequence satisfies the recurrence.”
Alfred van der Poorten — *A proof that Euler missed... (1979)*

* Someone's "sour comment" after Henri Cohen's report on Apéry's proof at the '78 ICM in Helsinki.

A victory for the French peasant...*

- The **Apéry numbers**

1, 5, 73, 1445, ...

satisfy

$$A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$$

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

THM
Apéry '78

$\zeta(3) = \sum_{n=1}^{\infty} \frac{1}{n^3}$ is irrational.



“After a few days of fruitless effort the specific problem was mentioned to Don Zagier (Bonn), and with *irritating speed* he showed that indeed the sequence satisfies the recurrence.”
Alfred van der Poorten — *A proof that Euler missed... (1979)*



Nowadays, there are excellent implementations of this **creative telescoping**, including:

- HolonomicFunctions** by Koutschan (Mathematica)
- Sigma** by Schneider (Mathematica)
- ore_algebra** by Kauers, Jaroschek, Johansson, Mezzarobba (Sage)

(These are just the ones I use on a regular basis...)

* Someone's "sour comment" after Henri Cohen's report on Apéry's proof at the '78 ICM in Helsinki.

Background: Creative telescoping

Goal

A telescoping recurrence for $\sum_{k=0}^n \underbrace{\binom{n}{k}^2 \binom{n+k}{k}^2}_{=: a(n,k)}$

N, K shift operators in n and k : $Na(n, k) = a(n+1, k)$



Marko Petkovsek, Herbert S. Wilf and Doron Zeilberger

$A = B$

A. K. Peters, Ltd., 1st edition, 1996

Background: Creative telescoping

Goal

A telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 =: a(n, k)$

N, K shift operators in n and k : $Na(n, k) = a(n+1, k)$

- Suppose we have $P(n, N) \in \mathbb{Q}[n, N]$ and $R(n, k) \in \mathbb{Q}(n, k)$ so that:

$$P(n, N)a(n, k) = (K - 1)R(n, k)a(n, k)$$



Marko Petkovsek, Herbert S. Wilf and Doron Zeilberger

$A = B$

A. K. Peters, Ltd., 1st edition, 1996

Background: Creative telescoping

Goal

A telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 =: a(n, k)$

N, K shift operators in n and k : $Na(n, k) = a(n+1, k)$

- Suppose we have $P(n, N) \in \mathbb{Q}[n, N]$ and $R(n, k) \in \mathbb{Q}(n, k)$ so that:

$$P(n, N)a(n, k) = (K - 1)R(n, k)a(n, k) = b(n, k+1) - b(n, k)$$



Marko Petkovsek, Herbert S. Wilf and Doron Zeilberger

$A = B$

A. K. Peters, Ltd., 1st edition, 1996

Background: Creative telescoping

Goal

A telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 =: a(n, k)$

N, K shift operators in n and k : $Na(n, k) = a(n+1, k)$

- Suppose we have $P(n, N) \in \mathbb{Q}[n, N]$ and $R(n, k) \in \mathbb{Q}(n, k)$ so that:

$$P(n, N)a(n, k) = (K - 1)R(n, k)a(n, k) = b(n, k+1) - b(n, k)$$

- Then: $P(n, N) \sum_{k \in \mathbb{Z}} a(n, k) = 0$

Assuming $\lim_{k \rightarrow \pm\infty} b(n, k) = 0$.



Marko Petkovsek, Herbert S. Wilf and Doron Zeilberger

$A = B$

A. K. Peters, Ltd., 1st edition, 1996

Background: Creative telescoping

Goal

A telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 =: a(n, k)$

N, K shift operators in n and k : $Na(n, k) = a(n+1, k)$

- Suppose we have $P(n, N) \in \mathbb{Q}[n, N]$ and $R(n, k) \in \mathbb{Q}(n, k)$ so that:

$$P(n, N)a(n, k) = (K - 1)R(n, k)a(n, k) = b(n, k+1) - b(n, k)$$

- Then: $P(n, N) \sum_{k \in \mathbb{Z}} a(n, k) = 0$

Assuming $\lim_{k \rightarrow \pm\infty} b(n, k) = 0$.

EG

$$P(n, N) = (n+2)^3 N^2 - (2n+3)(17n^2 + 51n + 39)N + (n+1)^3$$

$$R(n, k) = \frac{4k^4(2n+3)(4n^2 - 2k^2 + 12n + 3k + 8)}{(n-k+1)^2(n-k+2)^2}$$

$R(n, k)$ is the **certificate** of the telescoping recurrence operator $P(n, N)$.



Marko Petkovsek, Herbert S. Wilf and Doron Zeilberger

$A = B$

A. K. Peters, Ltd., 1st edition, 1996

Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order d :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order d :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

- If $\lim_{n \rightarrow \infty} p_k(n) = c_k$, then the **characteristic polynomial** is:

$$\lambda^d + c_{d-1} \lambda^{d-1} + \cdots + c_1 \lambda + c_0 = \prod_{k=1}^d (\lambda - \lambda_k)$$

Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order d :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

- If $\lim_{n \rightarrow \infty} p_k(n) = c_k$, then the **characteristic polynomial** is:

$$\lambda^d + c_{d-1} \lambda^{d-1} + \cdots + c_1 \lambda + c_0 = \prod_{k=1}^d (\lambda - \lambda_k)$$



THM
Poincaré
1885

Suppose the $|\lambda_k|$ are distinct. Then, for any solution u_n ,

$$\lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} = \lambda_k \quad (\text{P})$$

for some $k \in \{1, \dots, d\}$, unless u_n is eventually zero.

Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order d :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

- If $\lim_{n \rightarrow \infty} p_k(n) = c_k$, then the **characteristic polynomial** is:

$$\lambda^d + c_{d-1} \lambda^{d-1} + \cdots + c_1 \lambda + c_0 = \prod_{k=1}^d (\lambda - \lambda_k)$$



THM
Poincaré
1885

Suppose the $|\lambda_k|$ are distinct. Then, for any solution u_n ,

$$\lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} = \lambda_k$$


(P)

for some $k \in \{1, \dots, d\}$, unless u_n is eventually zero.

THM
Perron
1909

Suppose, in addition, $p_0(n) \neq 0$ for all $n \geq 0$.

Then, for each λ_k , there exists a u_n such that (P) holds.



Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order d :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

- If $\lim_{n \rightarrow \infty} p_k(n) = c_k$, then the **characteristic polynomial** is:

$$\lambda^d + c_{d-1} \lambda^{d-1} + \cdots + c_1 \lambda + c_0 = \prod_{k=1}^d (\lambda - \lambda_k)$$



THM
Poincaré
1885

Suppose the $|\lambda_k|$ are distinct. Then, for any solution u_n ,

$$\lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} = \lambda_k$$


(P)

for some $k \in \{1, \dots, d\}$, unless u_n is eventually zero.

THM
Perron
1909

Suppose, in addition, $p_0(n) \neq 0$ for all $n \geq 0$.

Then, for each λ_k , there exists a u_n such that (P) holds.



EG
Kooman
1989

For $u_{n+2} - 2u_{n+1} + (1 + \frac{1}{n^2})u_n = 0$, we have $\lambda_1 = \lambda_2 = 1$.
However, (P) does not hold for any real u_n .

There are two complex solutions asymptotic to n^r with $r = \exp(\pm\pi i/3)$.

Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order d :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

- If $\lim_{n \rightarrow \infty} p_k(n) = c_k$, then the **characteristic polynomial** is:

$$\lambda^d + c_{d-1} \lambda^{d-1} + \cdots + c_1 \lambda + c_0 = \prod_{k=1}^d (\lambda - \lambda_k)$$



THM
Poincaré
1885

Suppose the $|\lambda_k|$ are distinct. Then, for any solution u_n ,

$$\lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} = \lambda_k$$


(P)

for some $k \in \{1, \dots, d\}$, unless u_n is eventually zero.

THM
Perron
1909

Suppose, in addition, $p_0(n) \neq 0$ for all $n \geq 0$.

Then, for each λ_k , there exists a u_n such that (P) holds.



EG
Kooman
1989

For $\alpha_n u_{n+2} + (\alpha_{n+1} - \alpha_n) u_{n+1} - \alpha_{n+1} u_n = 0$, we have $\lambda_1, \lambda_2 = \pm 1$.

However, (P) holds for all u_n with RHS = 1.

$$\alpha_n = 1 + \frac{(-1)^n}{n}$$

Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n + 1)^3 u_{n+1} = (2n + 1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$: Apéry numbers $A(n)$ 1, 5, 73, 1445, 33001, \dots

Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n + 1)^3 u_{n+1} = (2n + 1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$: Apéry numbers $A(n)$

1, 5, 73, 1445, 33001, ...

- $u_0 = 0, u_1 = 1$: 2nd solution $B(n)$

0, 1, $\frac{117}{8}$, $\frac{62531}{216}$, $\frac{11424695}{1728}$, ...

Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$: Apéry numbers $A(n)$

1, 5, 73, 1445, 33001, ...

- $u_0 = 0, u_1 = 1$: 2nd solution $B(n)$

0, 1, $\frac{117}{8}$, $\frac{62531}{216}$, $\frac{11424695}{1728}$, ...

THM
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$: Apéry numbers $A(n)$ 1, 5, 73, 1445, 33001, \dots
- $u_0 = 0, u_1 = 1$: 2nd solution $B(n)$ 0, 1, $\frac{117}{8}$, $\frac{62531}{216}$, $\frac{11424695}{1728}$, \dots

THM
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

- Characteristic polynomial $n^2 - 34n + 1$ has roots $(1 \pm \sqrt{2})^4 \approx 33.97, 0.0294$.
 $A(n), B(n)$ grow like $(1 + \sqrt{2})^4$.

Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$: Apéry numbers $A(n)$ 1, 5, 73, 1445, 33001, \dots
- $u_0 = 0, u_1 = 1$: 2nd solution $B(n)$ 0, 1, $\frac{117}{8}$, $\frac{62531}{216}$, $\frac{11424695}{1728}$, \dots

THM
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

- Characteristic polynomial $n^2 - 34n + 1$ has roots $(1 \pm \sqrt{2})^4 \approx 33.97, 0.0294$.
 $A(n), B(n)$ grow like $(1 + \sqrt{2})^4$.
- By Perron's theorem, there is a (unique) solution

$$C(n) = \gamma A(n) + B(n) \quad \text{with} \quad \lim_{n \rightarrow \infty} \frac{C(n+1)}{C(n)} = (1 - \sqrt{2})^4.$$

Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$: Apéry numbers $A(n)$ 1, 5, 73, 1445, 33001, \dots
- $u_0 = 0, u_1 = 1$: 2nd solution $B(n)$ 0, 1, $\frac{117}{8}$, $\frac{62531}{216}$, $\frac{11424695}{1728}$, \dots

THM
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

- Characteristic polynomial $n^2 - 34n + 1$ has roots $(1 \pm \sqrt{2})^4 \approx 33.97, 0.0294$.
 $A(n), B(n)$ grow like $(1 + \sqrt{2})^4$.
- By Perron's theorem, there is a (unique) solution

$$C(n) = \gamma A(n) + B(n) \quad \text{with} \quad \lim_{n \rightarrow \infty} \frac{C(n+1)}{C(n)} = (1 - \sqrt{2})^4.$$
$$\downarrow$$
$$0 = \gamma + \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$$

Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$: Apéry numbers $A(n)$ 1, 5, 73, 1445, 33001, \dots
- $u_0 = 0, u_1 = 1$: 2nd solution $B(n)$ 0, 1, $\frac{117}{8}$, $\frac{62531}{216}$, $\frac{11424695}{1728}$, \dots

THM
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

- Characteristic polynomial $n^2 - 34n + 1$ has roots $(1 \pm \sqrt{2})^4 \approx 33.97, 0.0294$.
 $A(n), B(n)$ grow like $(1 + \sqrt{2})^4$.
- By Perron's theorem, there is a (unique) solution

$$C(n) = \gamma A(n) + B(n) \quad \text{with} \quad \lim_{n \rightarrow \infty} \frac{C(n+1)}{C(n)} = (1 - \sqrt{2})^4.$$

$$\downarrow$$

$$0 = \gamma + \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$$

COR $A(n)\zeta(3) - 6B(n)$ is "Perron's small solution".

This is a small linear form in 1 and $\zeta(3)$.

Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$: Apéry numbers $A(n)$ 1, 5, 73, 1445, 33001, \dots
- $u_0 = 0, u_1 = 1$: 2nd solution $B(n)$ 0, 1, $\frac{117}{8}$, $\frac{62531}{216}$, $\frac{11424695}{1728}$, \dots

THM
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

- Characteristic polynomial $n^2 - 34n + 1$ has roots $(1 \pm \sqrt{2})^4 \approx 33.97, 0.0294$.
 $A(n), B(n)$ grow like $(1 + \sqrt{2})^4$.
- By Perron's theorem, there is a (unique) solution

$$C(n) = \gamma A(n) + B(n) \quad \text{with} \quad \lim_{n \rightarrow \infty} \frac{C(n+1)}{C(n)} = (1 - \sqrt{2})^4.$$
$$\downarrow$$
$$0 = \gamma + \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$$

COR $A(n)\zeta(3) - 6B(n)$ is "Perron's small solution".

This is a small linear form in 1 and $\zeta(3)$.

? Tools to construct the solutions guaranteed by Perron's theorem?

A motivating example

- The (central) **Delannoy numbers** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}$ satisfy
 $(n+1)u_{n+1} = 3(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$
count lattice paths from $(0, 0)$ to (n, n) using the steps $(0, 1)$, $(1, 0)$ and $(1, 1)$

$$A(n) = 1, 3, 13, 63, 321, 1683, 8989, 48639, \dots$$



A motivating example

- The (central) **Delannoy numbers** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}$ satisfy
 $(n+1)u_{n+1} = 3(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$
count lattice paths from $(0, 0)$ to (n, n) using the steps $(0, 1)$, $(1, 0)$ and $(1, 1)$
- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 3, 13, 63, 321, 1683, 8989, 48639, \dots$$

$$B(n) = 0, 1, \frac{9}{2}, \frac{131}{6}, \frac{445}{4}, \frac{34997}{60}, \frac{62307}{20}, \frac{2359979}{140}, \dots$$



A motivating example

- The (central) **Delannoy numbers** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}$ satisfy
 $(n+1)u_{n+1} = 3(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$
count lattice paths from $(0, 0)$ to (n, n) using the steps $(0, 1)$, $(1, 0)$ and $(1, 1)$
- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 3, 13, 63, 321, 1683, 8989, 48639, \dots$$

$$B(n) = 0, 1, \frac{9}{2}, \frac{131}{6}, \frac{445}{4}, \frac{34997}{60}, \frac{62307}{20}, \frac{2359979}{140}, \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{3}, \frac{9}{26}, \frac{131}{378}, \frac{445}{1284}, \frac{34997}{100980}, \frac{62307}{179780}, \frac{2359979}{6809460}, \dots \rightarrow 0.34657359\dots$$



A motivating example

- The (central) **Delannoy numbers** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}$ satisfy
 $(n+1)u_{n+1} = 3(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$
count lattice paths from $(0, 0)$ to (n, n) using the steps $(0, 1)$, $(1, 0)$ and $(1, 1)$
- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 3, 13, 63, 321, 1683, 8989, 48639, \dots$$

$$B(n) = 0, 1, \frac{9}{2}, \frac{131}{6}, \frac{445}{4}, \frac{34997}{60}, \frac{62307}{20}, \frac{2359979}{140}, \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{3}, \frac{9}{26}, \frac{131}{378}, \frac{445}{1284}, \frac{34997}{100980}, \frac{62307}{179780}, \frac{2359979}{6809460}, \dots \rightarrow 0.34657359\dots$$

$$Q(n) - Q(n-1) = \frac{1}{3}, \frac{1}{78}, \frac{1}{2457}, \frac{1}{80892}, \frac{1}{2701215}, \frac{1}{90770922}, \frac{1}{3060511797}, \dots$$



A motivating example

- The (central) **Delannoy numbers** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}$ satisfy
 $(n+1)u_{n+1} = 3(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$
count lattice paths from $(0, 0)$ to (n, n) using the steps $(0, 1)$, $(1, 0)$ and $(1, 1)$
- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 3, 13, 63, 321, 1683, 8989, 48639, \dots$$

$$B(n) = 0, 1, \frac{9}{2}, \frac{131}{6}, \frac{445}{4}, \frac{34997}{60}, \frac{62307}{20}, \frac{2359979}{140}, \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{3}, \frac{9}{26}, \frac{131}{378}, \frac{445}{1284}, \frac{34997}{100980}, \frac{62307}{179780}, \frac{2359979}{6809460}, \dots \rightarrow 0.34657359\dots$$

$$Q(n) - Q(n-1) = \frac{1}{3}, \frac{1}{78}, \frac{1}{2457}, \frac{1}{80892}, \frac{1}{2701215}, \frac{1}{90770922}, \frac{1}{3060511797}, \dots$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{2} \ln 2$$



A motivating example

- The (central) **Delannoy polynomials** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$ satisfy $A(-1) = 0, A(0) = 1$
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$
- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) =$$

$$B(n) =$$

$$Q(n) := \frac{B(n)}{A(n)} =$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} =$$



A motivating example

- The (central) **Delannoy polynomials** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$ satisfy
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$
- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) =$$

$$Q(n) := \frac{B(n)}{A(n)} =$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} =$$



A motivating example

- The (central) **Delannoy polynomials** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$ satisfy
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$

- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} =$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} =$$



A motivating example

- The (central) **Delannoy polynomials** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$ satisfy
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$

- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1 + 2x}, \frac{3(1 + 2x)}{2(1 + 6x + 6x^2)}, \frac{11 + 60x + 60x^2}{6(1 + 2x)(1 + 10x + 10x^2)}, \dots$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} =$$



A motivating example

- The (central) **Delannoy polynomials** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$ satisfy
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$

- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1 + 2x}, \frac{3(1 + 2x)}{2(1 + 6x + 6x^2)}, \frac{11 + 60x + 60x^2}{6(1 + 2x)(1 + 10x + 10x^2)}, \dots$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} =$$

$$Q(3) = \frac{1}{2x} - \frac{1}{4x^2} + \frac{1}{6x^3} - \frac{1}{8x^4} + \frac{1}{10x^5} - \frac{1}{12x^6} + O(x^{-7})$$



A motivating example

- The (central) **Delannoy polynomials** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$ satisfy
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$

- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1 + 2x}, \frac{3(1 + 2x)}{2(1 + 6x + 6x^2)}, \frac{11 + 60x + 60x^2}{6(1 + 2x)(1 + 10x + 10x^2)}, \dots$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{2} \ln \left(1 + \frac{1}{x} \right)$$

$$Q(3) = \frac{1}{2x} - \frac{1}{4x^2} + \frac{1}{6x^3} - \frac{1}{8x^4} + \frac{1}{10x^5} - \frac{1}{12x^6} + O(x^{-7})$$



A motivating example

- The (central) **Delannoy polynomials** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$ satisfy
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$ $A(-1) = 0, A(0) = 1$

- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 1+2x, 1+6x+6x^2, (1+2x)(1+10x+10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1+2x), \frac{1}{6}(11+60x+60x^2), \frac{5}{12}(1+2x)(5+42x+42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1+2x}, \frac{3(1+2x)}{2(1+6x+6x^2)}, \frac{11+60x+60x^2}{6(1+2x)(1+10x+10x^2)}, \dots$$

$$Q(n) - Q(n-1) = \frac{1}{1+2x}, \frac{1}{2(1+2x)(1+6x+6x^2)}, \dots$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{2} \ln \left(1 + \frac{1}{x} \right)$$

$$Q(3) = \frac{1}{2x} - \frac{1}{4x^2} + \frac{1}{6x^3} - \frac{1}{8x^4} + \frac{1}{10x^5} - \frac{1}{12x^6} + O(x^{-7})$$



A motivating example

- The (central) **Delannoy polynomials** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$ satisfy $A(-1) = 0, A(0) = 1$
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$

- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1 + 2x}, \frac{3(1 + 2x)}{2(1 + 6x + 6x^2)}, \frac{11 + 60x + 60x^2}{6(1 + 2x)(1 + 10x + 10x^2)}, \dots$$

$$Q(n) - Q(n-1) = \frac{1}{1 + 2x}, \frac{1}{2(1 + 2x)(1 + 6x + 6x^2)}, \dots \quad \frac{1}{nA(n)A(n-1)}$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{2} \ln \left(1 + \frac{1}{x} \right)$$

$$Q(3) = \frac{1}{2x} - \frac{1}{4x^2} + \frac{1}{6x^3} - \frac{1}{8x^4} + \frac{1}{10x^5} - \frac{1}{12x^6} + O(x^{-7})$$



A motivating example

- The (central) **Delannoy polynomials** $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$ satisfy $A(-1) = 0, A(0) = 1$
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$

- Let $B(n)$ be the 2nd solution with initial conditions $B(0) = 0, B(1) = 1$.

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1 + 2x}, \frac{3(1 + 2x)}{2(1 + 6x + 6x^2)}, \frac{11 + 60x + 60x^2}{6(1 + 2x)(1 + 10x + 10x^2)}, \dots$$

$$Q(n) - Q(n-1) = \frac{1}{1 + 2x}, \frac{1}{2(1 + 2x)(1 + 6x + 6x^2)}, \dots \quad \frac{1}{nA(n)A(n-1)}$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{2} \ln \left(1 + \frac{1}{x} \right) = \sum_{n=1}^{\infty} \frac{1}{nA(n)A(n-1)}$$

$$Q(3) = \frac{1}{2x} - \frac{1}{4x^2} + \frac{1}{6x^3} - \frac{1}{8x^4} + \frac{1}{10x^5} - \frac{1}{12x^6} + O(x^{-7})$$



Explore Apéry limits

1 Pick a binomial sum $A(n)$.

Using creative telescoping, compute a recurrence satisfied by $A(n)$.

Explore Apéry limits

- 1 Pick a binomial sum $A(n)$.
Using creative telescoping, compute a recurrence satisfied by $A(n)$.
- 2 Compute the initial terms of a secondary solution $B(n)$ to the recurrence.

Explore Apéry limits

- 1 Pick a binomial sum $A(n)$.
Using creative telescoping, compute a recurrence satisfied by $A(n)$.
- 2 Compute the initial terms of a secondary solution $B(n)$ to the recurrence.
- 3 Try to identify $\lim_{n \rightarrow \infty} B(n)/A(n)$,
either numerically or as a power series in an additional parameter.

Explore Apéry limits

- 1 Pick a binomial sum $A(n)$.
Using creative telescoping, compute a recurrence satisfied by $A(n)$.
- 2 Compute the initial terms of a secondary solution $B(n)$ to the recurrence.
- 3 Try to identify $\lim_{n \rightarrow \infty} B(n)/A(n)$,
either numerically or as a power series in an additional parameter.

EG
HW

Use $\sum_{k=0}^n \binom{n}{k} \binom{n-k}{k} x^k$ to rediscover the CF

$$\arctan(z) = \frac{z}{1+} \frac{1^2 z^2}{3+} \frac{2^2 z^2}{5+} \cdots \frac{n^2 z^2}{(2n+1)+} \cdots$$

Explore Apéry limits

- 1 Pick a binomial sum $A(n)$.
Using creative telescoping, compute a recurrence satisfied by $A(n)$.
- 2 Compute the initial terms of a secondary solution $B(n)$ to the recurrence.
- 3 Try to identify $\lim_{n \rightarrow \infty} B(n)/A(n)$,
either numerically or as a power series in an additional parameter.

EG
HW

Use $\sum_{k=0}^n \binom{n}{k} \binom{n-k}{k} x^k$ to rediscover the CF

$$\arctan(z) = \frac{z}{1+} \frac{1^2 z^2}{3+} \frac{2^2 z^2}{5+} \cdots \frac{n^2 z^2}{(2n+1)+} \cdots$$

EG
HW

Start with $\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}^2 x^k$ and $\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}^3 x^k$.

Compare findings with those by Zudilin on simultaneous approximations to the logarithm, dilogarithm and trilogarithm.

Explore Apéry limits

- 1 Pick a binomial sum $A(n)$.
Using creative telescoping, compute a recurrence satisfied by $A(n)$.
- 2 Compute the initial terms of a secondary solution $B(n)$ to the recurrence.
- 3 Try to identify $\lim_{n \rightarrow \infty} B(n)/A(n)$,
either numerically or as a power series in an additional parameter.

EG
HW

Use $\sum_{k=0}^n \binom{n}{k} \binom{n-k}{k} x^k$ to rediscover the CF

$$\arctan(z) = \frac{z}{1+} \frac{1^2 z^2}{3+} \frac{2^2 z^2}{5+} \cdots \frac{n^2 z^2}{(2n+1)+} \cdots$$

EG
HW

Start with $\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}^2 x^k$ and $\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}^3 x^k$.

Compare findings with those by Zudilin on simultaneous approximations to the logarithm, dilogarithm and trilogarithm.

EG
bonus

For $\sum_{k=0}^n \binom{n}{k}^2 \binom{3k}{n}$, determine and prove the Apéry limits.

This is one of many cases conjectured by Almkvist, van Straten and Zudilin (2008) for CY DE's. Can we establish all these limits in a uniform fashion?

Approaches to proving Apéry limits

Q

How to prove $\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$?

1 Via explicit expressions:

(Apéry, '78)

$$B(n) = \frac{1}{6} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left(\sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right)$$

Approaches to proving Apéry limits

Q

How to prove $\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$?

1 Via explicit expressions:

(Apéry, '78)

$$B(n) = \frac{1}{6} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left(\sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right)$$

2 Via integral representations:

(Beukers, '79)

$$(-1)^n \int_0^1 \int_0^1 \int_0^1 \frac{x^n (1-x)^n y^n (1-y)^n z^n (1-z)^n}{(1 - (1-xy)z)^{n+1}} dx dy dz = A(n)\zeta(3) - 6B(n)$$

Approaches to proving Apéry limits

Q

How to prove $\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$?

1 Via explicit expressions:

(Apéry, '78)

$$B(n) = \frac{1}{6} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left(\sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right)$$

2 Via integral representations:

(Beukers, '79)

$$(-1)^n \int_0^1 \int_0^1 \int_0^1 \frac{x^n (1-x)^n y^n (1-y)^n z^n (1-z)^n}{(1 - (1-xy)z)^{n+1}} dx dy dz = A(n)\zeta(3) - 6B(n)$$

3 Via hypergeometric series representations:

(Gutnik, '79)

$$-\frac{1}{2} \sum_{t=1}^{\infty} R'_n(t) = A(n)\zeta(3) - 6B(n), \quad \text{where } R_n(t) = \left(\frac{(t-1) \cdots (t-n)}{t(t+1) \cdots (t+n)} \right)^2$$

Approaches to proving Apéry limits

Q

How to prove $\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$?

1 Via explicit expressions:

(Apéry, '78)

$$B(n) = \frac{1}{6} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left(\sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right)$$

2 Via integral representations:

(Beukers, '79)

$$(-1)^n \int_0^1 \int_0^1 \int_0^1 \frac{x^n (1-x)^n y^n (1-y)^n z^n (1-z)^n}{(1 - (1-xy)z)^{n+1}} dx dy dz = A(n)\zeta(3) - 6B(n)$$

3 Via hypergeometric series representations:

(Gutnik, '79)

$$-\frac{1}{2} \sum_{t=1}^{\infty} R'_n(t) = A(n)\zeta(3) - 6B(n), \quad \text{where } R_n(t) = \left(\frac{(t-1) \cdots (t-n)}{t(t+1) \cdots (t+n)} \right)^2$$

4 Via modular forms

(Beukers '87, Zagier '03, Yang '07)

Approaches to proving Apéry limits

Q

How to prove $\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$?

1 Via explicit expressions:

(Apéry, '78)

$$B(n) = \frac{1}{6} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left(\sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right)$$

2 Via integral representations:

(Beukers, '79)

$$(-1)^n \int_0^1 \int_0^1 \int_0^1 \frac{x^n (1-x)^n y^n (1-y)^n z^n (1-z)^n}{(1 - (1-xy)z)^{n+1}} dx dy dz = A(n)\zeta(3) - 6B(n)$$

3 Via hypergeometric series representations:

(Gutnik, '79)

$$-\frac{1}{2} \sum_{t=1}^{\infty} R'_n(t) = A(n)\zeta(3) - 6B(n), \quad \text{where } R_n(t) = \left(\frac{(t-1) \cdots (t-n)}{t(t+1) \cdots (t+n)} \right)^2$$

4 Via modular forms

(Beukers '87, Zagier '03, Yang '07)

5 Via continued fractions (for recurrences of order 2)

Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 +} \frac{a_2}{b_2 +} \frac{a_3}{b_3 +} \dots := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

THM 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit: $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

THM 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit: $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

- Here, $A(n), B(n)$ are the solutions to $u_n = b_n u_{n-1} + a_n u_{n-2}$
with $A(-1) = 0, A(0) = 1$ and $B(0) = 0, B(1) = a_1$.

Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

THM 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit: $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

- Here, $A(n), B(n)$ are the solutions to $u_n = b_n u_{n-1} + a_n u_{n-2}$
with $A(-1) = 0, A(0) = 1$ and $B(0) = 0, B(1) = a_1$.

proof The n -th convergent is $C_n := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \dots \frac{a_n}{b_n}}} = \frac{B(n)}{A(n)}$. □

Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

THM 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit: $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

- Here, $A(n), B(n)$ are the solutions to $u_n = b_n u_{n-1} + a_n u_{n-2}$
with $A(-1) = 0, A(0) = 1$ and $B(0) = 0, B(1) = a_1$.

EG

$$A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k \text{ solves } (n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}.$$

Let $B(n)$ be the solution with $B(0) = 0, B(1) = 1$.

Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

THM 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit: $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

- Here, $A(n), B(n)$ are the solutions to $u_n = b_n u_{n-1} + a_n u_{n-2}$
with $A(-1) = 0, A(0) = 1$ and $B(0) = 0, B(1) = a_1$.

EG

$$A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k \text{ solves } (n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}.$$

Let $B(n)$ be the solution with $B(0) = 0, B(1) = 1$.

$$\text{Hence, } n!A(n), n!B(n) \text{ solve } u_{n+1} = \frac{(2x+1)(2n+1)}{b_{n+1}} u_n - \frac{n^2}{a_{n+1}} u_{n-1}.$$

Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

THM 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit: $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

- Here, $A(n), B(n)$ are the solutions to $u_n = b_n u_{n-1} + a_n u_{n-2}$
with $A(-1) = 0, A(0) = 1$ and $B(0) = 0, B(1) = a_1$.

EG

$$A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k \text{ solves } (n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}.$$

Let $B(n)$ be the solution with $B(0) = 0, B(1) = 1$.

$$\text{Hence, } n!A(n), n!B(n) \text{ solve } u_{n+1} = \frac{(2x+1)(2n+1)}{b_{n+1}} u_n - \frac{n^2}{a_{n+1}} u_{n-1}.$$


Apéry limit and equivalent CF:

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{(2x+1) - \frac{1^2}{3(2x+1) - \frac{2^2}{5(2x+1) - \dots}}} = \frac{1}{2} \ln \left(1 + \frac{1}{x} \right)$$

Franel numbers

DEF
Franel
1894

$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$ are the (generalized) Franel numbers.




Franel numbers

DEF
Franel
1894

$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$ are the (generalized) Franel numbers.

- $A^{(1)}(n) = 2^n$

$$u_{n+1} = 2u_n$$



Franel numbers

DEF
Franel
1894


$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$ are the (generalized) Franel numbers.

- $A^{(1)}(n) = 2^n$

$$u_{n+1} = 2u_n$$

- $A^{(2)}(n) = \binom{2n}{n}$

$$(n+1)u_{n+1} = 2(2n+1)u_n$$



Franel numbers

DEF
Franel
1894

$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$ are the (generalized) Franel numbers.

- $A^{(1)}(n) = 2^n$


$$u_{n+1} = 2u_n$$

- $A^{(2)}(n) = \binom{2n}{n}$

$$(n+1)u_{n+1} = 2(2n+1)u_n$$

- $A^{(3)}(n) = 1, 2, 10, 56, 346, 2252, 15184, 104960, 739162, \dots$

$$(n+1)^2 u_{n+1} = (7n^2 + 7n + 2)u_n + 8n^2 u_{n-1}$$



(Franel, 1894)

Franel numbers

DEF
Franel
1894

$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$ are the (generalized) Franel numbers.

- $A^{(1)}(n) = 2^n$

$$u_{n+1} = 2u_n$$

- $A^{(2)}(n) = \binom{2n}{n}$

$$(n+1)u_{n+1} = 2(2n+1)u_n$$

- $A^{(3)}(n) = 1, 2, 10, 56, 346, 2252, 15184, 104960, 739162, \dots$


$$(n+1)^2 u_{n+1} = (7n^2 + 7n + 2)u_n + 8n^2 u_{n-1}$$

(Franel, 1894)

- $A^{(4)}(n) = 1, 2, 18, 164, 1810, 21252, 263844, 3395016, 44916498, \dots$

$$(n+1)^3 u_{n+1} = 2(2n+1)(3n^2 + 3n + 1)u_n + 4n(16n^2 - 1)u_{n-1}$$

(Franel, 1895)



Franel numbers

DEF
Franel
1894

$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$ are the (generalized) Franel numbers.

- $A^{(1)}(n) = 2^n$

$$u_{n+1} = 2u_n$$

- $A^{(2)}(n) = \binom{2n}{n}$

$$(n+1)u_{n+1} = 2(2n+1)u_n$$

- $A^{(3)}(n) = 1, 2, 10, 56, 346, 2252, 15184, 104960, 739162, \dots$


$$(n+1)^2 u_{n+1} = (7n^2 + 7n + 2)u_n + 8n^2 u_{n-1}$$

(Franel, 1894)

- $A^{(4)}(n) = 1, 2, 18, 164, 1810, 21252, 263844, 3395016, 44916498, \dots$

$$(n+1)^3 u_{n+1} = 2(2n+1)(3n^2 + 3n + 1)u_n + 4n(16n^2 - 1)u_{n-1}$$

(Franel, 1895)



CONJ
Franel,
1895

The minimal recurrence for $A^{(s)}(n)$ has order $\lfloor \frac{s+1}{2} \rfloor$
and degree $s - 1$.

(spoiler: the degree part is not true)

Frenel's conjecture

CONJ

Frenel,
1895

The minimal recurrence for $A^{(s)}(n)$ has order $\lfloor \frac{s+1}{2} \rfloor$ and degree $s - 1$.

Franel's conjecture

CONJ
Franel,
1895

The minimal recurrence for $A^{(s)}(n)$ has order $\lfloor \frac{s+1}{2} \rfloor$ and degree $s - 1$.

- Perlstadt '86: order 3 recurrences for $s = 5, 6$ of degrees 6, 9 computed using MACSYMA and creative telescoping



Franel's conjecture

CONJ
Franel,
1895

The minimal recurrence for $A^{(s)}(n)$ has order $\lfloor \frac{s+1}{2} \rfloor$ and degree $s - 1$.

- Perlstadt '86: order 3 recurrences for $s = 5, 6$ of degrees 6, 9
computed using MACSYMA and creative telescoping

THM
Stoll '97

$A^{(s)}(n)$ satisfies a recurrence of order $\lfloor \frac{s+1}{2} \rfloor$.

Cusick '89 also constructs such recurrences.



Franel's conjecture

CONJ
Franel,
1895

The minimal recurrence for $A^{(s)}(n)$ has order $\lfloor \frac{s+1}{2} \rfloor$ and degree $s - 1$.

- Perlstadt '86: order 3 recurrences for $s = 5, 6$ of degrees 6, 9 computed using MACSYMA and creative telescoping



THM
Stoll '97

$A^{(s)}(n)$ satisfies a recurrence of order $\lfloor \frac{s+1}{2} \rfloor$.

Cusick '89 also constructs such recurrences.



CONJ
Bostan
'21

The minimal recurrence for $A^{(s)}(n)$ has order $m = \lfloor \frac{s+1}{2} \rfloor$ and

$$\text{degree} = \begin{cases} \frac{1}{3}m(m^2 - 1) + 1, & \text{for even } s, \\ \frac{1}{3}m^3 - \frac{1}{2}m^2 + \frac{2}{3}m + \frac{(-1)^{m-1}}{4}, & \text{for odd } s. \end{cases}$$

If true, the degree grows like $s^3/24$.



Franel's conjecture

CONJ
Franel,
1895

The minimal recurrence for $A^{(s)}(n)$ has order $\lfloor \frac{s+1}{2} \rfloor$ and degree $s - 1$.

- Perlstadt '86: order 3 recurrences for $s = 5, 6$ of degrees 6, 9 computed using MACSYMA and creative telescoping

THM
Stoll '97

$A^{(s)}(n)$ satisfies a recurrence of order $\lfloor \frac{s+1}{2} \rfloor$.

Cusick '89 also constructs such recurrences.

CONJ
Bostan
'21

The minimal recurrence for $A^{(s)}(n)$ has order $m = \lfloor \frac{s+1}{2} \rfloor$ and

$$\text{degree} = \begin{cases} \frac{1}{3}m(m^2 - 1) + 1, & \text{for even } s, \\ \frac{1}{3}m^3 - \frac{1}{2}m^2 + \frac{2}{3}m + \frac{(-1)^m - 1}{4}, & \text{for odd } s. \end{cases}$$

If true, the degree grows like $s^3/24$.

- Verified at least for $s \leq 20$.
using MinimalRecurrence from the LREtools Maple package



Franel's conjecture

CONJ

Franel,
1895

The minimal recurrence for $A^{(s)}(n)$ has order $\lfloor \frac{s+1}{2} \rfloor$ and degree $s - 1$.

- Perlstadt '86: order 3 recurrences for $s = 5, 6$ of degrees 6, 9 computed using MACSYMA and creative telescoping



THM

Stoll '97

$A^{(s)}(n)$ satisfies a recurrence of order $\lfloor \frac{s+1}{2} \rfloor$.

Cusick '89 also constructs such recurrences.



CONJ

Bostan
'21

The minimal recurrence for $A^{(s)}(n)$ has order $m = \lfloor \frac{s+1}{2} \rfloor$ and

$$\text{degree} = \begin{cases} \frac{1}{3}m(m^2 - 1) + 1, & \text{for even } s, \\ \frac{1}{3}m^3 - \frac{1}{2}m^2 + \frac{2}{3}m + \frac{(-1)^{m-1}}{4}, & \text{for odd } s. \end{cases}$$

If true, the degree grows like $s^3/24$.



- Verified at least for $s \leq 20$.
using MinimalRecurrence from the LREtools Maple package
- Goal: The minimal **telescoping** recurrence for $A^{(s)}(n)$ has order $\geq \lfloor \frac{s+1}{2} \rfloor$.

How to prove lower bounds for orders of recurrences?

EG

- $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$: recurrence of order 2 (Apéry '78)

- $\sum_{k=0}^n \binom{n}{k}^s$: recurrence of order $\lfloor \frac{s+1}{2} \rfloor$ (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

How to prove lower bounds for orders of recurrences?

EG

- $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$: recurrence of order 2 (Apéry '78)

- $\sum_{k=0}^n \binom{n}{k}^s$: recurrence of order $\lfloor \frac{s+1}{2} \rfloor$ (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

- For fixed sequence, order 1 can be ruled out using Hyper, (Petkovšek '92)
an algorithm to compute order 1 (right) factors of recurrence operators.

How to prove lower bounds for orders of recurrences?

EG

- $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$: recurrence of order 2 (Apéry '78)

- $\sum_{k=0}^n \binom{n}{k}^s$: recurrence of order $\lfloor \frac{s+1}{2} \rfloor$ (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

- For fixed sequence, order 1 can be ruled out using Hyper, (Petkovšek '92)
an algorithm to compute order 1 (right) factors of recurrence operators.
- For Franel numbers, order 1 can be ruled out for all $s \geq 3$ (Yuan-Lu-Schmidt '08)
using congruential properties.

How to prove lower bounds for orders of recurrences?

EG

- $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$: recurrence of order 2 (Apéry '78)

- $\sum_{k=0}^n \binom{n}{k}^s$: recurrence of order $\lfloor \frac{s+1}{2} \rfloor$ (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

- For fixed sequence, order 1 can be ruled out using Hyper, (Petkovšek '92)
an algorithm to compute order 1 (right) factors of recurrence operators.
- For Franel numbers, order 1 can be ruled out for all $s \geq 3$ (Yuan-Lu-Schmidt '08)
using congruential properties.
- There are algorithms for fixed recurrence operators (Beke 1894, Bronstein '94,
for computing factors of differen(tial/ce) operators. Zhou-van Hoeij '19, ...)

How to prove lower bounds for orders of recurrences?

EG

• $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$: recurrence of order 2 (Apéry '78)

• $\sum_{k=0}^n \binom{n}{k}^s$: recurrence of order $\lfloor \frac{s+1}{2} \rfloor$ (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

- For fixed sequence, order 1 can be ruled out using Hyper, (Petkovšek '92)
an algorithm to compute order 1 (right) factors of recurrence operators.
- For Franel numbers, order 1 can be ruled out for all $s \geq 3$ (Yuan-Lu-Schmidt '08)
using congruential properties.
- There are algorithms for fixed recurrence operators (Beke 1894, Bronstein '94,
for computing factors of differen(tial/ce) operators. Zhou-van Hoeij '19, ...)
- If $A(n+1)/A(n) \rightarrow \mu$ for $\mu \in \bar{\mathbb{Q}}$ of degree d , then $A(n)$ cannot satisfy a
recurrence over \mathbb{Q} of order less than d . (McIntosh '89)

How to prove lower bounds for orders of recurrences?

EG

- $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$: recurrence of order 2 (Apéry '78)

- $\sum_{k=0}^n \binom{n}{k}^s$: recurrence of order $\lfloor \frac{s+1}{2} \rfloor$ (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

- For fixed sequence, order 1 can be ruled out using Hyper, (Petkovšek '92)
an algorithm to compute order 1 (right) factors of recurrence operators.
- For Franel numbers, order 1 can be ruled out for all $s \geq 3$ (Yuan-Lu-Schmidt '08)
using congruential properties.
- There are algorithms for fixed recurrence operators (Beke 1894, Bronstein '94,
for computing factors of differen(tial/ce) operators. Zhou-van Hoeij '19, ...)
- If $A(n+1)/A(n) \rightarrow \mu$ for $\mu \in \bar{\mathbb{Q}}$ of degree d , then $A(n)$ cannot satisfy a
recurrence over \mathbb{Q} of order less than d . (McIntosh '89)

For Apéry numbers: $\mu = (1 + \sqrt{2})^4$.

For Franel numbers: $\mu = 2^s$. Not helpful!

Solutions to the Franel number recurrences

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

Solutions to the Franel number recurrences

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof
outline

- Suppose: $P(n, N) \binom{n}{k}^s = b(n, k+1) - b(n, k)$
for a hypergeometric term $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$.



Solutions to the Franel number recurrences

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof
outline

- Suppose: $P(n, N) \binom{n}{k-t}^s = b(n, k-t+1) - b(n, k-t)$
for a hypergeometric term $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$.



Solutions to the Franel number recurrences

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof
outline

- 1 Suppose: $P(n, N) \binom{n}{k-t}^s = b(n, k-t+1) - b(n, k-t)$
for a hypergeometric term $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$.
- 2 $P(n, N) \sum_{k=\alpha}^{\beta-1} \binom{n}{k-t}^s = b(n, \beta-t) - b(n, \alpha-t)$ $b(n, t)$ entire
for large n



Solutions to the Franel number recurrences

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof
outline

- 1 Suppose: $P(n, N) \binom{n}{k-t}^s = b(n, k-t+1) - b(n, k-t)$
for a hypergeometric term $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$.
- 2 $P(n, N) \sum_{k=\alpha}^{\beta-1} \binom{n}{k-t}^s = b(n, \beta-t) - b(n, \alpha-t)$ $b(n, t)$ entire
for large n
- 3 $P(n, N) \sum_{k \in \mathbb{Z}} \binom{n}{k-t}^s = 0$ since $b(n, t) = \text{rat}(n, t) \binom{n}{t}^s$ and
 $\binom{n}{t} = O\left(\frac{1}{t^{n+1}}\right)$ for real $t \rightarrow \pm\infty$

□

Solutions to the Franel number recurrences

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof
outline

1 Suppose: $P(n, N) \binom{n}{k-t}^s = b(n, k-t+1) - b(n, k-t)$
for a hypergeometric term $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$.

2 $P(n, N) \sum_{k=\alpha}^{\beta-1} \binom{n}{k-t}^s = b(n, \beta-t) - b(n, \alpha-t)$ $b(n, t)$ entire
for large n

3 $P(n, N) \sum_{k \in \mathbb{Z}} \binom{n}{k-t}^s = 0$ since $b(n, t) = \text{rat}(n, t) \binom{n}{t}^s$ and
 $\binom{n}{t} = O\left(\frac{1}{t^{n+1}}\right)$ for real $t \rightarrow \pm\infty$

4 $A^{(s)}(n, t) = \left(\frac{\pi t}{\sin(\pi t)}\right)^s \sum_{k=0}^n \binom{n}{k-t}^s$

□

Solutions to the Franel number recurrences

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof
outline

1 Suppose: $P(n, N) \binom{n}{k-t}^s = b(n, k-t+1) - b(n, k-t)$
for a hypergeometric term $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$.

2 $P(n, N) \sum_{k=\alpha}^{\beta-1} \binom{n}{k-t}^s = b(n, \beta-t) - b(n, \alpha-t)$ $b(n, t)$ entire
for large n

3 $P(n, N) \sum_{k \in \mathbb{Z}} \binom{n}{k-t}^s = 0$ since $b(n, t) = \text{rat}(n, t) \binom{n}{t}^s$ and
 $\binom{n}{t} = O\left(\frac{1}{t^{n+1}}\right)$ for real $t \rightarrow \pm\infty$

4 $A^{(s)}(n, t) = \left(\frac{\pi t}{\sin(\pi t)}\right)^s \sum_{k=0}^n \binom{n}{k-t}^s = \left(\frac{\pi t}{\sin(\pi t)}\right)^s \sum_{k \in \mathbb{Z}} \binom{n}{k-t}^s + O(t^s)$

5 Hence, $P(n, N)A^{(s)}(n, t) = O(t^s)$. □

Apéry limits for Franel numbers

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

Apéry limits for Franel numbers

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

THM
S-Zudilin
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Apéry limits for Franel numbers

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

THM
S-Zudilin
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

- This follows from locally uniform convergence in t of

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s}}{\sum_{k=0}^n \binom{n}{k}^s} = \left(\frac{\pi t}{\sin(\pi t)} \right)^s.$$

Apéry limits for Franel numbers

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

THM
S-Zudilin
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

- This follows from locally uniform convergence in t of

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=0}^n \binom{n}{k}^s \left[\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s}}{\sum_{k=0}^n \binom{n}{k}^s} = \left(\frac{\pi t}{\sin(\pi t)} \right)^s.$$

- For large n and $k \approx n/2$,

$$\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \approx \prod_{j=1}^{\infty} \left(1 - \frac{t}{j}\right) \left(1 + \frac{t}{j}\right) = \frac{\sin(\pi t)}{\pi t}.$$

Apéry limits for Franel numbers

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

THM
S-Zudilin
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

$$\left(\frac{\pi t}{\sin(\pi t)} \right)^s = \left(\sum_{j=1}^{\infty} \left(2 - \frac{1}{2^{2j-2}} \right) \zeta(2j) t^{2j} \right)^s = 1 + s\zeta(2) t^2 + \frac{s(5s+2)}{4} \zeta(4) t^4 + O(t^6)$$

Apéry limits for Franel numbers

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

THM
S-Zudilin
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

$$\left(\frac{\pi t}{\sin(\pi t)} \right)^s = \left(\sum_{j=1}^{\infty} \left(2 - \frac{1}{2^{2j-2}} \right) \zeta(2j) t^{2j} \right)^s = 1 + s\zeta(2) t^2 + \frac{s(5s+2)}{4} \zeta(4) t^4 + O(t^6)$$

EG
 $j=1$

$$\lim_{n \rightarrow \infty} \frac{B^{(s)}(n)}{A^{(s)}(n)} = \frac{1}{s(s+1)} s\zeta(2) = \frac{\zeta(2)}{s+1}$$

$$B^{(s)}(n) = \frac{A_1^{(s)}(n)}{A_1^{(s)}(1)} = 0, 1, \dots$$

$$A_1^{(s)}(1) = s(s+1)$$

Apéry limits for Franel numbers

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

THM
S-Zudilin
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

$$\left(\frac{\pi t}{\sin(\pi t)} \right)^s = \left(\sum_{j=1}^{\infty} \left(2 - \frac{1}{2^{2j-2}} \right) \zeta(2j) t^{2j} \right)^s = 1 + s\zeta(2) t^2 + \frac{s(5s+2)}{4} \zeta(4) t^4 + O(t^6)$$

EG
 $j=1$

$$\lim_{n \rightarrow \infty} \frac{B^{(s)}(n)}{A^{(s)}(n)} = \frac{1}{s(s+1)} s\zeta(2) = \frac{\zeta(2)}{s+1}$$

$$B^{(s)}(n) = \frac{A_1^{(s)}(n)}{A_1^{(s)}(1)} = 0, 1, \dots$$

- $s = 3, 4$ numerically observed by Cusick (1979)
- $s = 3$ proved by Zagier (2009)
- $s = 5$ conjectured by Almkvist, van Straten, Zudilin (2008)
- $s \geq 3$ conjectured by Chamberland-S (2020)

$$A_1^{(s)}(1) = s(s+1)$$

Apéry limits for Franel numbers

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

THM
S-Zudilin
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

$$\left(\frac{\pi t}{\sin(\pi t)} \right)^s = \left(\sum_{j=1}^{\infty} \left(2 - \frac{1}{2^{2j-2}} \right) \zeta(2j) t^{2j} \right)^s = 1 + s\zeta(2) t^2 + \frac{s(5s+2)}{4} \zeta(4) t^4 + O(t^6)$$

EG
 $j = 1$

$$\lim_{n \rightarrow \infty} \frac{B^{(s)}(n)}{A^{(s)}(n)} = \frac{1}{s(s+1)} s\zeta(2) = \frac{\zeta(2)}{s+1}$$

$$B^{(s)}(n) = \frac{A_1^{(s)}(n)}{A_1^{(s)}(1)} = 0, 1, \dots$$

- $s = 3, 4$ numerically observed by Cusick (1979)
- $s = 3$ proved by Zagier (2009)
- $s = 5$ conjectured by Almkvist, van Straten, Zudilin (2008)
- $s \geq 3$ conjectured by Chamberland-S (2020)

$$A_1^{(s)}(1) = s(s+1)$$

EG
 $j = 2$

$$\lim_{n \rightarrow \infty} \frac{C^{(s)}(n)}{A^{(s)}(n)} = \frac{12}{s(s+1)(s+2)(s+3)} \frac{s(5s+2)}{4} \zeta(4) \quad C^{(s)}(n) = \frac{A_2^{(s)}(n)}{A_2^{(s)}(1)} = 0, 1, \dots$$

- $s \geq 5$ conjectured by Chamberland-S (2020)

Telescoping version of Franel's conjecture

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ has order at least $\lfloor \frac{s+1}{2} \rfloor$.

Telescoping version of Franel's conjecture

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for $s \leq 30$.

Telescoping version of Franel's conjecture

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for $s \leq 30$.
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

Telescoping version of Franel's conjecture

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for $s \leq 30$.
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by $A_j^{(s)}(n) \in \mathbb{Q}$ if $0 \leq 2j < s$.

Here, and below, we assume that n is large enough.



Telescoping version of Franel's conjecture

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for $s \leq 30$.
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by $A_j^{(s)}(n) \in \mathbb{Q}$ if $0 \leq 2j < s$.

Here, and below, we assume that n is large enough.

2 The claim follows if these are linearly independent.



Telescoping version of Franel's conjecture

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for $s \leq 30$.
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by $A_j^{(s)}(n) \in \mathbb{Q}$ if $0 \leq 2j < s$.

Here, and below, we assume that n is large enough.

2 The claim follows if these are linearly independent.

3 $0 = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j A_j^{(s)}(n)$

$\lambda_j \in \mathbb{Q}$

□

Telescoping version of Franel's conjecture

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for $s \leq 30$.
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by $A_j^{(s)}(n) \in \mathbb{Q}$ if $0 \leq 2j < s$.

Here, and below, we assume that n is large enough.

2 The claim follows if these are linearly independent.

$$3 \quad 0 = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j A_j^{(s)}(n) \quad \implies \quad 0 = \lim_{n \rightarrow \infty} \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j \frac{A_j^{(s)}(n)}{A^{(s)}(n)}$$

$\lambda_j \in \mathbb{Q}$

□

Telescoping version of Franel's conjecture

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for $s \leq 30$.
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by $A_j^{(s)}(n) \in \mathbb{Q}$ if $0 \leq 2j < s$.

Here, and below, we assume that n is large enough.

2 The claim follows if these are linearly independent.

$$3 \quad 0 = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j A_j^{(s)}(n) \quad \implies \quad 0 = \lim_{n \rightarrow \infty} \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j \varphi_j \pi^{2j}$$

$\lambda_j \in \mathbb{Q}$ $\varphi_j \in \mathbb{Q}^\times$

□

Telescoping version of Franel's conjecture

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for $s \leq 30$.
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by $A_j^{(s)}(n) \in \mathbb{Q}$ if $0 \leq 2j < s$.

Here, and below, we assume that n is large enough.

2 The claim follows if these are linearly independent.

$$3 \quad 0 = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j A_j^{(s)}(n) \quad \implies \quad 0 = \lim_{n \rightarrow \infty} \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j \varphi_j \pi^{2j}$$

$\lambda_j \in \mathbb{Q}$ $\varphi_j \in \mathbb{Q}^\times$

4 Transcendence of π implies that all λ_j are zero. □

When does creative telescoping fall short?

EG
Paule,
Schorn
'95

$$\text{Consider } S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n}.$$



When does creative telescoping fall short?

EG
Paule,
Schorn
'95

Consider $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n}$.

- Any telescoping recurrence $P(n, N)$ has order $\geq d - 1$:

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$



When does creative telescoping fall short?

EG
Paule,
Schorn
'95

Consider $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$.

- Any telescoping recurrence $P(n, N)$ has order $\geq d - 1$:

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However, $S_d(n) = (-d)^n$ satisfies $NS_d(n) + dS_d(n) = 0$.



When does creative telescoping fall short?

EG
Paule,
Schorn
'95

Consider $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$.

- Any telescoping recurrence $P(n, N)$ has order $\geq d - 1$:

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However, $S_d(n) = (-d)^n$ satisfies $NS_d(n) + dS_d(n) = 0$.



- Open problem: When does CT fall short?

When does creative telescoping fall short?

EG
Paule,
Schorn
'95

Consider $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$.

- Any telescoping recurrence $P(n, N)$ has order $\geq d - 1$:

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However, $S_d(n) = (-d)^n$ satisfies $NS_d(n) + dS_d(n) = 0$.



- Open problem: When does CT fall short?
- Can these cases be “fixed” by a different hypergeometric representation?

When does creative telescoping fall short?

EG
Paule,
Schorn
'95

Consider $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$.

- Any telescoping recurrence $P(n, N)$ has order $\geq d - 1$:

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However, $S_d(n) = (-d)^n$ satisfies $NS_d(n) + dS_d(n) = 0$.



- Open problem: When does CT fall short?
- Can these cases be “fixed” by a different hypergeometric representation?

EG
Riese '01
“creative
symmetrizing”

Consider $\sum_{k=1}^{2n} (-1)^k \binom{2n}{k}^2 \binom{2n}{k-1} = (-1)^n \frac{(3n)!}{n!^2(n-1)!(2n+1)}$.



When does creative telescoping fall short?

EG
Paule,
Schorn
'95

Consider $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$.

- Any telescoping recurrence $P(n, N)$ has order $\geq d - 1$:

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However, $S_d(n) = (-d)^n$ satisfies $NS_d(n) + dS_d(n) = 0$.



- Open problem: When does CT fall short?
- Can these cases be “fixed” by a different hypergeometric representation?

EG
Riese '01
“creative
symmetrizing”

Consider $\sum_{k=1}^{2n} (-1)^k \binom{2n}{k}^2 \binom{2n}{k-1} = (-1)^n \frac{(3n)!}{n!^2(n-1)!(2n+1)}$.

- CT produces order 2 recurrence on summand $a(n, k)$, but



When does creative telescoping fall short?

EG
Paule,
Schorn
'95

Consider $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$.

- Any telescoping recurrence $P(n, N)$ has order $\geq d - 1$:

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However, $S_d(n) = (-d)^n$ satisfies $NS_d(n) + dS_d(n) = 0$.



- Open problem: When does CT fall short?
- Can these cases be “fixed” by a different hypergeometric representation?

EG
Riese '01
“creative
symmetrizing”

Consider $\sum_{k=1}^{2n} (-1)^k \binom{2n}{k}^2 \binom{2n}{k-1} = (-1)^n \frac{(3n)!}{n!^2(n-1)!(2n+1)}$.

- CT produces order 2 recurrence on summand $a(n, k)$, but
- order 1 on $a(n, k) + a(n, 2n - k + 1) = \frac{2n - 2k + 1}{2n - k + 1} a(n, k)$.



When does creative telescoping fall short?

EG
Paule,
Schorn
'95

Consider $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$.

- Any telescoping recurrence $P(n, N)$ has order $\geq d - 1$:

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However, $S_d(n) = (-d)^n$ satisfies $NS_d(n) + dS_d(n) = 0$.



- Open problem: When does CT fall short?
- Can these cases be “fixed” by a different hypergeometric representation?

EG
Riese '01
“creative
symmetrizing”

Consider $\sum_{k=1}^{2n} (-1)^k \binom{2n}{k}^2 \binom{2n}{k-1} = (-1)^n \frac{(3n)!}{n!^2(n-1)!(2n+1)}$.

- CT produces order 2 recurrence on summand $a(n, k)$, but
- order 1 on $a(n, k) + a(n, 2n - k + 1) = \frac{2n - 2k + 1}{2n - k + 1} a(n, k)$.



“ Studying a huge number of practical applications one is tempted to conjecture that Zeilberger’s algorithm always returns the recurrence with minimal order. ”
Peter Paule, Markus Schorn, *Journal of Symbolic Computation*, 1995

Modularity and Apéry limits

- Let $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$ be Zagier's sporadic sequence **C**.

1, 3, 15, 93, ...

THM
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left(\frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4)$$

$$x(\tau) = q - 4q^2 + 10q^3 + O(q^4)$$

$$q = e^{2\pi i \tau}$$

Modularity and Apéry limits

- Let $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$ be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

THM
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left(\frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4)$$

$$x(\tau) = q - 4q^2 + 10q^3 + O(q^4)$$

$$q = e^{2\pi i \tau}$$

- Context:
 - $f(\tau)$ modular form of weight k
 - $x(\tau)$ modular function
 - $y(x)$ such that $y(x(\tau)) = f(\tau)$

Then $y(x)$ satisfies a linear differential equation $Ly = 0$ of order $k + 1$.

Modularity and Apéry limits

- Let $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$ be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

THM
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left(\frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4)$$

$$x(\tau) = q - 4q^2 + 10q^3 + O(q^4)$$

$$q = e^{2\pi i \tau}$$

- Context:
 - $f(\tau)$ modular form of weight k
 - $x(\tau)$ modular function
 - $y(x)$ such that $y(x(\tau)) = f(\tau)$

Then $y(x)$ satisfies a linear differential equation $Ly = 0$ of order $k + 1$.

- Solutions to $Ly = \text{rat}(x)$ are of the form $y(x)$ times an Eichler integral of

$$h(\tau) = \left(\frac{Dx(\tau)}{x(\tau)} \right)^{k+1} \frac{\text{rat}(x(\tau))}{f(\tau)} \quad (\text{a modular form of weight } k+2) \quad (\text{Yang '07})$$

$$D = q \frac{d}{dq}$$

If $\sum_{n \geq 1} c_n q^n$ is a modular form of weight $k + 2$, then $\sum_{n \geq 1} \frac{c_n}{n^{k+1}} q^n$ is an Eichler integral.

Modularity and Apéry limits

- Let $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$ be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

THM
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left(\frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4)$$

$$x(\tau) = q - 4q^2 + 10q^3 + O(q^4)$$

$$q = e^{2\pi i \tau}$$

- $F(x) := \sum_{n \geq 0} A(n)x^n \implies F(x(\tau)) = f(\tau)$
- $G(x) := \sum_{n \geq 0} B(n)x^n \implies G(x(\tau)) = f(\tau) \sum_{n \geq 1} \frac{\binom{-3}{n}}{n^2} \frac{q^n}{1+q^n}$

Modularity and Apéry limits

- Let $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$ be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

THM
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left(\frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4)$$

$$x(\tau) = q - 4q^2 + 10q^3 + O(q^4)$$

$$q = e^{2\pi i \tau}$$

- $F(x) := \sum_{n \geq 0} A(n)x^n \implies F(x(\tau)) = f(\tau)$
- $G(x) := \sum_{n \geq 0} B(n)x^n \implies G(x(\tau)) = f(\tau) \sum_{n \geq 1} \frac{\binom{-3}{n}}{n^2} \frac{q^n}{1+q^n}$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \lim_{x \rightarrow \frac{1}{9}} \frac{G(x)}{F(x)}$$

characteristic roots 1, 9

$F(x), G(x)$ have radius of convergence $R = \frac{1}{9}$.

$G(x) - LF(x)$ has radius of convergence $R = 1 > \frac{1}{9}$ for $L = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$.

Modularity and Apéry limits

- Let $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$ be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

THM
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left(\frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4) \quad x(\tau) = q - 4q^2 + 10q^3 + O(q^4) \quad q = e^{2\pi i}$$

- $F(x) := \sum_{n \geq 0} A(n)x^n \implies F(x(\tau)) = f(\tau)$
- $G(x) := \sum_{n \geq 0} B(n)x^n \implies G(x(\tau)) = f(\tau) \sum_{n \geq 1} \frac{\binom{-3}{n}}{n^2} \frac{q^n}{1+q^n}$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \lim_{x \rightarrow \frac{1}{9}} \frac{G(x)}{F(x)} = \lim_{\tau \rightarrow 0} \frac{G(x(\tau))}{F(x(\tau))}$$

characteristic roots 1, 9

$$x(\tau) = \frac{1}{9} \text{ for } \tau = 0 \text{ or } q = 1$$

$F(x), G(x)$ have radius of convergence $R = \frac{1}{9}$.

$G(x) - LF(x)$ has radius of convergence $R = 1 > \frac{1}{9}$ for $L = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$.

Modularity and Apéry limits

- Let $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$ be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

THM
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left(\frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4) \quad x(\tau) = q - 4q^2 + 10q^3 + O(q^4) \quad q = e^{2\pi i}$$

- $F(x) := \sum_{n \geq 0} A(n)x^n \implies F(x(\tau)) = f(\tau)$
- $G(x) := \sum_{n \geq 0} B(n)x^n \implies G(x(\tau)) = f(\tau) \sum_{n \geq 1} \frac{\binom{-3}{n}}{n^2} \frac{q^n}{1+q^n}$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \lim_{x \rightarrow \frac{1}{9}} \frac{G(x)}{F(x)} = \lim_{\tau \rightarrow 0} \frac{G(x(\tau))}{F(x(\tau))} = \lim_{q \rightarrow 1} \sum_{n \geq 1} \frac{\binom{-3}{n}}{n^2} \frac{q^n}{1+q^n} = \frac{1}{2} L_{-3}(2)$$

characteristic roots 1, 9

$$x(\tau) = \frac{1}{9} \text{ for } \tau = 0 \text{ or } q = 1$$

$F(x), G(x)$ have radius of convergence $R = \frac{1}{9}$.

$G(x) - LF(x)$ has radius of convergence $R = 1 > \frac{1}{9}$ for $L = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$.

Conclusions and some open questions

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover, $A_j^{(s)}(n)$ with $0 \leq 2j < s$ are linearly independent, so that any telescoping recurrence has order at least $\lfloor \frac{s+1}{2} \rfloor$.

Conclusions and some open questions

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover, $A_j^{(s)}(n)$ with $0 \leq 2j < s$ are linearly independent, so that any telescoping recurrence has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- Cusick '89 and Stoll '97 construct recurrences for Franel numbers. Can these constructions produce telescoping recurrences?

Conclusions and some open questions

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover, $A_j^{(s)}(n)$ with $0 \leq 2j < s$ are linearly independent, so that any telescoping recurrence has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- Cusick '89 and Stoll '97 construct recurrences for Franel numbers.
Can these constructions produce telescoping recurrences?
- What can we learn from other families of binomial sums?
In particular, it would be nice to simplify some of the technical steps in the arguments.

Conclusions and some open questions

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover, $A_j^{(s)}(n)$ with $0 \leq 2j < s$ are linearly independent, so that any telescoping recurrence has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- Cusick '89 and Stoll '97 construct recurrences for Franel numbers.
Can these constructions produce telescoping recurrences?
- What can we learn from other families of binomial sums?
In particular, it would be nice to simplify some of the technical steps in the arguments.
- Can we (uniformly) establish the conjectural Apéry limits for CY DE's?

Conclusions and some open questions

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover, $A_j^{(s)}(n)$ with $0 \leq 2j < s$ are linearly independent, so that any telescoping recurrence has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- Cusick '89 and Stoll '97 construct recurrences for Franel numbers.
Can these constructions produce telescoping recurrences?
- What can we learn from other families of binomial sums?
In particular, it would be nice to simplify some of the technical steps in the arguments.
- Can we (uniformly) establish the conjectural Apéry limits for CY DE's?
- How to explicitly construct the solutions guaranteed by Perron's theorem?

Conclusions and some open questions

THM
S-Zudilin
'21

Any telescoping recurrence for $\sum_{k=0}^n \binom{n}{k}^s$ solved by $A_j^{(s)}(n)$ if $0 \leq 2j < s$.
(fine print: for large enough n)

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left(\frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover, $A_j^{(s)}(n)$ with $0 \leq 2j < s$ are linearly independent, so that any telescoping recurrence has order at least $\lfloor \frac{s+1}{2} \rfloor$.

- Cusick '89 and Stoll '97 construct recurrences for Franel numbers.
Can these constructions produce telescoping recurrences?
- What can we learn from other families of binomial sums?
In particular, it would be nice to simplify some of the technical steps in the arguments.
- Can we (uniformly) establish the conjectural Apéry limits for CY DE's?
- How to explicitly construct the solutions guaranteed by Perron's theorem?
- Can we explain when CT falls short? And algorithmically "fix" this issue?

THANK YOU!

Slides for this talk will be available from my website:
<http://arminstraub.com/talks>



M. Chamberland, A. Straub

Apéry limits: Experiments and proofs

American Mathematical Monthly, Vol. 128, Nr. 9, 2021, p. 811-824



A. Straub, W. Zudilin

Sums of powers of binomials, their Apéry limits, and Franel's suspicions

arXiv:2112.09576