

# Sums of powers of binomials, their Apéry limits, and Franel's suspicions

Joint Seminar: MATHEXP-PoISys & Transcendence and Combinatorics  
Inria Saclay & Sorbonne University


Armin Straub

July 1, 2022

University of South Alabama

**CONJ**  $\pi, \zeta(3), \zeta(5), \dots$  are algebraically independent over  $\mathbb{Q}$ .

- Apéry (1978):  $\zeta(3)$  is irrational
- Open:  $\zeta(5)$  is irrational
- Open:  $\zeta(3)$  is transcendental
- Open:  $\zeta(3)/\pi^3$  is irrational



based on joint work(s) with:



Marc Chamberland  
(Grinnell College)




Wadim Zudilin  
(Radboud University)

# The last shall be first: conclusions

**CONJ**  
Franel,  
1895

The minimal recurrence for  $A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$  has order  $\lfloor \frac{s+1}{2} \rfloor$ .



**THM**  
Stoll '97


$A^{(s)}(n)$  satisfies a recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$ .



# The last shall be first: conclusions

**CONJ**  
Frenel,  
1895

The minimal recurrence for  $A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$  has order  $\lfloor \frac{s+1}{2} \rfloor$ .



**THM**  
Stoll '97

$A^{(s)}(n)$  satisfies a recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$ .



$$\sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$


**THM**  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

# The last shall be first: conclusions

**CONJ**  
Franel,  
1895

The minimal recurrence for  $A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$  has order  $\lfloor \frac{s+1}{2} \rfloor$ .



**THM**  
Stoll '97

$A^{(s)}(n)$  satisfies a recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$ .



$$\sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

**THM**  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover,  $A_j^{(s)}(n)$  with  $0 \leq 2j < s$  are linearly independent, so that any telescoping recurrence has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

# Apéry numbers and the irrationality of $\zeta(3)$

- The **Apéry numbers**

1, 5, 73, 1445, ...

satisfy

$$A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$$

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

**THM**  
Apéry '78

$\zeta(3) = \sum_{n=1}^{\infty} \frac{1}{n^3}$  is irrational.



# Apéry numbers and the irrationality of $\zeta(3)$

- The **Apéry numbers**

1, 5, 73, 1445, ...

satisfy

$$A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$$

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$



**THM**  
Apéry '78

$\zeta(3) = \sum_{n=1}^{\infty} \frac{1}{n^3}$  is irrational.

**proof** The same recurrence is satisfied by the “near”-integers

$$B(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left( \sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right).$$

Then,  $\frac{B(n)}{A(n)} \rightarrow \zeta(3)$ . But too fast for  $\zeta(3)$  to be rational.  $\square$

# Apéry numbers and the irrationality of $\zeta(3)$

- The **Apéry numbers**

1, 5, 73, 1445, ...

satisfy

$$A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$$

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

**THM**  
Apéry '78

$\zeta(3) = \sum_{n=1}^{\infty} \frac{1}{n^3}$  is irrational.



“After a few days of fruitless effort the specific problem was mentioned to Don Zagier (Bonn), and with *irritating speed* he showed that indeed the sequence satisfies the recurrence.”  
**Alfred van der Poorten** — *A proof that Euler missed...* (1979)



# Apéry numbers and the irrationality of $\zeta(3)$

- The **Apéry numbers**

1, 5, 73, 1445, ...

satisfy

$$A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$$

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

**THM**  
Apéry '78

$\zeta(3) = \sum_{n=1}^{\infty} \frac{1}{n^3}$  is irrational.



“After a few days of fruitless effort the specific problem was mentioned to Don Zagier (Bonn), and with *irritating speed* he showed that indeed the sequence satisfies the recurrence.”  
**Alfred van der Poorten** — *A proof that Euler missed...* (1979)



Nowadays, there are many excellent implementations of this **creative telescoping**.



# Background: Creative telescoping

## Goal

A telescoping recurrence for  $\sum_{k=0}^n \underbrace{\binom{n}{k}^2 \binom{n+k}{k}^2}_{=: a(n,k)}$

$N, K$  shift operators in  $n$  and  $k$ :  $Na(n, k) = a(n+1, k)$



Marko Petkovsek, Herbert S. Wilf and Doron Zeilberger

$A = B$

A. K. Peters, Ltd., 1st edition, 1996

# Background: Creative telescoping

## Goal

A telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 =: a(n, k)$

$N, K$  shift operators in  $n$  and  $k$ :  $Na(n, k) = a(n+1, k)$

- Suppose we have  $P(n, N) \in \mathbb{Q}[n, N]$  and  $R(n, k) \in \mathbb{Q}(n, k)$  so that:

$$P(n, N)a(n, k) = (K - 1)R(n, k)a(n, k)$$



Marko Petkovsek, Herbert S. Wilf and Doron Zeilberger

$A = B$

A. K. Peters, Ltd., 1st edition, 1996

# Background: Creative telescoping

## Goal

A telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 =: a(n, k)$

$N, K$  shift operators in  $n$  and  $k$ :  $Na(n, k) = a(n+1, k)$

- Suppose we have  $P(n, N) \in \mathbb{Q}[n, N]$  and  $R(n, k) \in \mathbb{Q}(n, k)$  so that:

$$P(n, N)a(n, k) = (K - 1)R(n, k)a(n, k) = b(n, k+1) - b(n, k)$$



Marko Petkovsek, Herbert S. Wilf and Doron Zeilberger

$A = B$

A. K. Peters, Ltd., 1st edition, 1996

# Background: Creative telescoping

## Goal

A telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 =: a(n, k)$

$N, K$  shift operators in  $n$  and  $k$ :  $Na(n, k) = a(n+1, k)$

- Suppose we have  $P(n, N) \in \mathbb{Q}[n, N]$  and  $R(n, k) \in \mathbb{Q}(n, k)$  so that:

$$P(n, N)a(n, k) = (K - 1)R(n, k)a(n, k) = b(n, k+1) - b(n, k)$$

- Then:  $P(n, N) \sum_{k \in \mathbb{Z}} a(n, k) = 0$

Assuming  $\lim_{k \rightarrow \pm\infty} b(n, k) = 0$ .



Marko Petkovsek, Herbert S. Wilf and Doron Zeilberger

$A = B$

A. K. Peters, Ltd., 1st edition, 1996

# Background: Creative telescoping

## Goal

A telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 =: a(n, k)$

$N, K$  shift operators in  $n$  and  $k$ :  $Na(n, k) = a(n+1, k)$

- Suppose we have  $P(n, N) \in \mathbb{Q}[n, N]$  and  $R(n, k) \in \mathbb{Q}(n, k)$  so that:

$$P(n, N)a(n, k) = (K - 1)R(n, k)a(n, k) = b(n, k+1) - b(n, k)$$

- Then:  $P(n, N) \sum_{k \in \mathbb{Z}} a(n, k) = 0$

Assuming  $\lim_{k \rightarrow \pm\infty} b(n, k) = 0$ .

## EG

$$P(n, N) = (n+2)^3 N^2 - (2n+3)(17n^2 + 51n + 39)N + (n+1)^3$$

$$R(n, k) = \frac{4k^4(2n+3)(4n^2 - 2k^2 + 12n + 3k + 8)}{(n-k+1)^2(n-k+2)^2}$$

$R(n, k)$  is the **certificate** of the telescoping recurrence operator  $P(n, N)$ .



Marko Petkovsek, Herbert S. Wilf and Doron Zeilberger

$A = B$

A. K. Peters, Ltd., 1st edition, 1996

## Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order  $d$ :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

## Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order  $d$ :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

- If  $\lim_{n \rightarrow \infty} p_k(n) = c_k$ , then the **characteristic polynomial** is:

$$\lambda^d + c_{d-1} \lambda^{d-1} + \cdots + c_1 \lambda + c_0 = \prod_{k=1}^d (\lambda - \lambda_k)$$

# Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order  $d$ :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

- If  $\lim_{n \rightarrow \infty} p_k(n) = c_k$ , then the **characteristic polynomial** is:

$$\lambda^d + c_{d-1} \lambda^{d-1} + \cdots + c_1 \lambda + c_0 = \prod_{k=1}^d (\lambda - \lambda_k)$$



**THM**  
Poincaré  
1885

Suppose the  $|\lambda_k|$  are distinct. Then, for any solution  $u_n$ ,

$$\lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} = \lambda_k \quad (\text{P})$$

for some  $k \in \{1, \dots, d\}$ , unless  $u_n$  is eventually zero.



# Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order  $d$ :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

- If  $\lim_{n \rightarrow \infty} p_k(n) = c_k$ , then the **characteristic polynomial** is:

$$\lambda^d + c_{d-1} \lambda^{d-1} + \cdots + c_1 \lambda + c_0 = \prod_{k=1}^d (\lambda - \lambda_k)$$



**THM**  
Poincaré  
1885

Suppose the  $|\lambda_k|$  are distinct. Then, for any solution  $u_n$ ,

$$\lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} = \lambda_k$$


(P)

for some  $k \in \{1, \dots, d\}$ , unless  $u_n$  is eventually zero.

**THM**  
Perron  
1909

Suppose, in addition,  $p_0(n) \neq 0$  for all  $n \geq 0$ .

Then, for each  $\lambda_k$ , there exists a  $u_n$  such that (P) holds.



# Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order  $d$ :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

- If  $\lim_{n \rightarrow \infty} p_k(n) = c_k$ , then the **characteristic polynomial** is:

$$\lambda^d + c_{d-1} \lambda^{d-1} + \cdots + c_1 \lambda + c_0 = \prod_{k=1}^d (\lambda - \lambda_k)$$



**THM**  
Poincaré  
1885

Suppose the  $|\lambda_k|$  are distinct. Then, for any solution  $u_n$ ,

$$\lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} = \lambda_k$$


(P)

for some  $k \in \{1, \dots, d\}$ , unless  $u_n$  is eventually zero.

**THM**  
Perron  
1909

Suppose, in addition,  $p_0(n) \neq 0$  for all  $n \geq 0$ .

Then, for each  $\lambda_k$ , there exists a  $u_n$  such that (P) holds.



**EG**  
Kooman  
1989

For  $u_{n+2} - 2u_{n+1} + (1 + \frac{1}{n^2})u_n = 0$ , we have  $\lambda_1 = \lambda_2 = 1$ .  
However, (P) does not hold for any real  $u_n$ .

There are two complex solutions asymptotic to  $n^r$  with  $r = \exp(\pm\pi i/3)$ .

# Background: Poincaré and Perron

- Normalized general homogeneous linear recurrence of order  $d$ :

$$u_{n+d} + p_{d-1}(n) u_{n+d-1} + \cdots + p_1(n) u_{n+1} + p_0(n) u_n = 0$$

- If  $\lim_{n \rightarrow \infty} p_k(n) = c_k$ , then the **characteristic polynomial** is:

$$\lambda^d + c_{d-1} \lambda^{d-1} + \cdots + c_1 \lambda + c_0 = \prod_{k=1}^d (\lambda - \lambda_k)$$



**THM**  
Poincaré  
1885

Suppose the  $|\lambda_k|$  are distinct. Then, for any solution  $u_n$ ,

$$\lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} = \lambda_k$$


(P)

for some  $k \in \{1, \dots, d\}$ , unless  $u_n$  is eventually zero.

**THM**  
Perron  
1909

Suppose, in addition,  $p_0(n) \neq 0$  for all  $n \geq 0$ .

Then, for each  $\lambda_k$ , there exists a  $u_n$  such that (P) holds.



**EG**  
Kooman  
1989

For  $\alpha_n u_{n+2} + (\alpha_{n+1} - \alpha_n) u_{n+1} - \alpha_{n+1} u_n = 0$ , we have  $\lambda_1, \lambda_2 = \pm 1$ .

However, (P) holds for all  $u_n$  with RHS = 1.

$$\alpha_n = 1 + \frac{(-1)^n}{n}$$

## Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n + 1)^3 u_{n+1} = (2n + 1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$ : Apéry numbers  $A(n)$  1, 5, 73, 1445, 33001, \dots

## Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n + 1)^3 u_{n+1} = (2n + 1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$ : Apéry numbers  $A(n)$

1, 5, 73, 1445, 33001, ...

- $u_0 = 0, u_1 = 1$ : 2nd solution  $B(n)$

0, 1,  $\frac{117}{8}$ ,  $\frac{62531}{216}$ ,  $\frac{11424695}{1728}$ , ...

## Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$ : Apéry numbers  $A(n)$

1, 5, 73, 1445, 33001, ...

- $u_0 = 0, u_1 = 1$ : 2nd solution  $B(n)$

0, 1,  $\frac{117}{8}$ ,  $\frac{62531}{216}$ ,  $\frac{11424695}{1728}$ , ...

**THM**  
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

## Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$ : Apéry numbers  $A(n)$  1, 5, 73, 1445, 33001, \dots
- $u_0 = 0, u_1 = 1$ : 2nd solution  $B(n)$  0, 1,  $\frac{117}{8}$ ,  $\frac{62531}{216}$ ,  $\frac{11424695}{1728}$ , \dots

**THM**  
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

- Characteristic polynomial  $n^2 - 34n + 1$  has roots  $(1 \pm \sqrt{2})^4 \approx 33.97, 0.0294$ .  
 $A(n), B(n)$  grow like  $(1 + \sqrt{2})^{4n}$ .

## Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$ : Apéry numbers  $A(n)$  1, 5, 73, 1445, 33001, \dots
- $u_0 = 0, u_1 = 1$ : 2nd solution  $B(n)$  0, 1,  $\frac{117}{8}$ ,  $\frac{62531}{216}$ ,  $\frac{11424695}{1728}$ , \dots

**THM**  
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

- Characteristic polynomial  $n^2 - 34n + 1$  has roots  $(1 \pm \sqrt{2})^4 \approx 33.97, 0.0294$ .  
 $A(n), B(n)$  grow like  $(1 + \sqrt{2})^{4n}$ .
- By Perron's theorem, there is a (unique) solution

$$C(n) = \gamma A(n) + B(n) \quad \text{with} \quad \lim_{n \rightarrow \infty} \frac{C(n+1)}{C(n)} = (1 - \sqrt{2})^4.$$



## Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$ : Apéry numbers  $A(n)$  1, 5, 73, 1445, 33001, \dots
- $u_0 = 0, u_1 = 1$ : 2nd solution  $B(n)$  0, 1,  $\frac{117}{8}$ ,  $\frac{62531}{216}$ ,  $\frac{11424695}{1728}$ , \dots

**THM**  
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

- Characteristic polynomial  $n^2 - 34n + 1$  has roots  $(1 \pm \sqrt{2})^4 \approx 33.97, 0.0294$ .  
 $A(n), B(n)$  grow like  $(1 + \sqrt{2})^{4n}$ .
- By Perron's theorem, there is a (unique) solution

$$C(n) = \gamma A(n) + B(n) \quad \text{with} \quad \lim_{n \rightarrow \infty} \frac{C(n+1)}{C(n)} = (1 - \sqrt{2})^4.$$
$$\downarrow$$
$$0 = \gamma + \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$$

## Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$ : Apéry numbers  $A(n)$  1, 5, 73, 1445, 33001, \dots
- $u_0 = 0, u_1 = 1$ : 2nd solution  $B(n)$  0, 1,  $\frac{117}{8}$ ,  $\frac{62531}{216}$ ,  $\frac{11424695}{1728}$ , \dots

**THM**  
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

- Characteristic polynomial  $n^2 - 34n + 1$  has roots  $(1 \pm \sqrt{2})^4 \approx 33.97, 0.0294$ .  
 $A(n), B(n)$  grow like  $(1 + \sqrt{2})^{4n}$ .
- By Perron's theorem, there is a (unique) solution

$$C(n) = \gamma A(n) + B(n) \quad \text{with} \quad \lim_{n \rightarrow \infty} \frac{C(n+1)}{C(n)} = (1 - \sqrt{2})^4.$$
$$\downarrow$$
$$0 = \gamma + \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$$

**COR**  $A(n)\zeta(3) - 6B(n)$  is "Perron's small solution".

This is a small linear form in 1 and  $\zeta(3)$ .

## Another look at Apéry's recurrence and limit

- Apéry's recurrence has order 2 and degree 3:

$$(n+1)^3 u_{n+1} = (2n+1)(17n^2 + 17n + 5)u_n - n^3 u_{n-1}.$$

- $u_{-1} = 0, u_0 = 1$ : Apéry numbers  $A(n)$  1, 5, 73, 1445, 33001, \dots
- $u_0 = 0, u_1 = 1$ : 2nd solution  $B(n)$  0, 1,  $\frac{117}{8}$ ,  $\frac{62531}{216}$ ,  $\frac{11424695}{1728}$ , \dots

**THM**  
Apéry '78

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$$

- Characteristic polynomial  $n^2 - 34n + 1$  has roots  $(1 \pm \sqrt{2})^4 \approx 33.97, 0.0294$ .  
 $A(n), B(n)$  grow like  $(1 + \sqrt{2})^{4n}$ .
- By Perron's theorem, there is a (unique) solution

$$C(n) = \gamma A(n) + B(n) \quad \text{with} \quad \lim_{n \rightarrow \infty} \frac{C(n+1)}{C(n)} = (1 - \sqrt{2})^4.$$
$$\downarrow$$
$$0 = \gamma + \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$$

**COR**  $A(n)\zeta(3) - 6B(n)$  is "Perron's small solution".

This is a small linear form in 1 and  $\zeta(3)$ .

**?** Tools to construct the solutions guaranteed by Perron's theorem?

# A motivating example

- The (central) **Delannoy numbers**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}$  satisfy  
 $(n+1)u_{n+1} = 3(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$   
count lattice paths from  $(0, 0)$  to  $(n, n)$  using the steps  $(0, 1)$ ,  $(1, 0)$  and  $(1, 1)$

$$A(n) = 1, 3, 13, 63, 321, 1683, 8989, 48639, \dots$$



# A motivating example

- The (central) **Delannoy numbers**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}$  satisfy  
 $(n+1)u_{n+1} = 3(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$   
count lattice paths from  $(0, 0)$  to  $(n, n)$  using the steps  $(0, 1)$ ,  $(1, 0)$  and  $(1, 1)$
- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 3, 13, 63, 321, 1683, 8989, 48639, \dots$$

$$B(n) = 0, 1, \frac{9}{2}, \frac{131}{6}, \frac{445}{4}, \frac{34997}{60}, \frac{62307}{20}, \frac{2359979}{140}, \dots$$



# A motivating example

- The (central) **Delannoy numbers**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}$  satisfy  
 $(n+1)u_{n+1} = 3(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$   
count lattice paths from  $(0, 0)$  to  $(n, n)$  using the steps  $(0, 1)$ ,  $(1, 0)$  and  $(1, 1)$
- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 3, 13, 63, 321, 1683, 8989, 48639, \dots$$

$$B(n) = 0, 1, \frac{9}{2}, \frac{131}{6}, \frac{445}{4}, \frac{34997}{60}, \frac{62307}{20}, \frac{2359979}{140}, \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{3}, \frac{9}{26}, \frac{131}{378}, \frac{445}{1284}, \frac{34997}{100980}, \frac{62307}{179780}, \frac{2359979}{6809460}, \dots \rightarrow 0.34657359\dots$$



# A motivating example

- The (central) **Delannoy numbers**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}$  satisfy  
 $(n+1)u_{n+1} = 3(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$   
count lattice paths from  $(0, 0)$  to  $(n, n)$  using the steps  $(0, 1)$ ,  $(1, 0)$  and  $(1, 1)$
- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 3, 13, 63, 321, 1683, 8989, 48639, \dots$$

$$B(n) = 0, 1, \frac{9}{2}, \frac{131}{6}, \frac{445}{4}, \frac{34997}{60}, \frac{62307}{20}, \frac{2359979}{140}, \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{3}, \frac{9}{26}, \frac{131}{378}, \frac{445}{1284}, \frac{34997}{100980}, \frac{62307}{179780}, \frac{2359979}{6809460}, \dots \rightarrow 0.34657359\dots$$

$$Q(n) - Q(n-1) = \frac{1}{3}, \frac{1}{78}, \frac{1}{2457}, \frac{1}{80892}, \frac{1}{2701215}, \frac{1}{90770922}, \frac{1}{3060511797}, \dots$$



# A motivating example

- The (central) **Delannoy numbers**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}$  satisfy  
 $(n+1)u_{n+1} = 3(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$   
count lattice paths from  $(0, 0)$  to  $(n, n)$  using the steps  $(0, 1)$ ,  $(1, 0)$  and  $(1, 1)$
- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 3, 13, 63, 321, 1683, 8989, 48639, \dots$$

$$B(n) = 0, 1, \frac{9}{2}, \frac{131}{6}, \frac{445}{4}, \frac{34997}{60}, \frac{62307}{20}, \frac{2359979}{140}, \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{3}, \frac{9}{26}, \frac{131}{378}, \frac{445}{1284}, \frac{34997}{100980}, \frac{62307}{179780}, \frac{2359979}{6809460}, \dots \rightarrow 0.34657359\dots$$

$$Q(n) - Q(n-1) = \frac{1}{3}, \frac{1}{78}, \frac{1}{2457}, \frac{1}{80892}, \frac{1}{2701215}, \frac{1}{90770922}, \frac{1}{3060511797}, \dots$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{2} \ln 2$$





# A motivating example

- The (central) **Delannoy polynomials**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$  satisfy  
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$
- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) =$$

$$B(n) =$$

$$Q(n) := \frac{B(n)}{A(n)} =$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} =$$



# A motivating example

- The (central) **Delannoy polynomials**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$  satisfy  
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$
- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) =$$

$$Q(n) := \frac{B(n)}{A(n)} =$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} =$$



# A motivating example

- The (central) **Delannoy polynomials**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$  satisfy  
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$
- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} =$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} =$$



# A motivating example

- The (central) **Delannoy polynomials**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$  satisfy  
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$

- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1 + 2x}, \frac{3(1 + 2x)}{2(1 + 6x + 6x^2)}, \frac{11 + 60x + 60x^2}{6(1 + 2x)(1 + 10x + 10x^2)}, \dots$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} =$$



# A motivating example

- The (central) **Delannoy polynomials**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$  satisfy  
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$

- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1 + 2x}, \frac{3(1 + 2x)}{2(1 + 6x + 6x^2)}, \frac{11 + 60x + 60x^2}{6(1 + 2x)(1 + 10x + 10x^2)}, \dots$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} =$$

$$Q(3) = \frac{1}{2x} - \frac{1}{4x^2} + \frac{1}{6x^3} - \frac{1}{8x^4} + \frac{1}{10x^5} - \frac{1}{12x^6} + O(x^{-7})$$



# A motivating example

- The (central) **Delannoy polynomials**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$  satisfy  
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$   $A(-1) = 0, A(0) = 1$

- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1 + 2x}, \frac{3(1 + 2x)}{2(1 + 6x + 6x^2)}, \frac{11 + 60x + 60x^2}{6(1 + 2x)(1 + 10x + 10x^2)}, \dots$$

$$Q(n) - Q(n-1) =$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{2} \ln \left( 1 + \frac{1}{x} \right)$$

$$Q(3) = \frac{1}{2x} - \frac{1}{4x^2} + \frac{1}{6x^3} - \frac{1}{8x^4} + \frac{1}{10x^5} - \frac{1}{12x^6} + O(x^{-7})$$



# A motivating example

- The (central) **Delannoy polynomials**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$  satisfy  $A(-1) = 0, A(0) = 1$   
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$

- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 1+2x, 1+6x+6x^2, (1+2x)(1+10x+10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1+2x), \frac{1}{6}(11+60x+60x^2), \frac{5}{12}(1+2x)(5+42x+42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1+2x}, \frac{3(1+2x)}{2(1+6x+6x^2)}, \frac{11+60x+60x^2}{6(1+2x)(1+10x+10x^2)}, \dots$$

$$Q(n) - Q(n-1) = \frac{1}{1+2x}, \frac{1}{2(1+2x)(1+6x+6x^2)}, \dots$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{2} \ln \left( 1 + \frac{1}{x} \right)$$

$$Q(3) = \frac{1}{2x} - \frac{1}{4x^2} + \frac{1}{6x^3} - \frac{1}{8x^4} + \frac{1}{10x^5} - \frac{1}{12x^6} + O(x^{-7})$$



# A motivating example

- The (central) **Delannoy polynomials**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$  satisfy  $A(-1) = 0, A(0) = 1$   
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$

- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1+2x}, \frac{3(1+2x)}{2(1+6x+6x^2)}, \frac{11+60x+60x^2}{6(1+2x)(1+10x+10x^2)}, \dots$$

$$Q(n) - Q(n-1) = \frac{1}{1+2x}, \frac{1}{2(1+2x)(1+6x+6x^2)}, \dots \quad \frac{1}{nA(n)A(n-1)}$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{2} \ln \left( 1 + \frac{1}{x} \right)$$

$$Q(3) = \frac{1}{2x} - \frac{1}{4x^2} + \frac{1}{6x^3} - \frac{1}{8x^4} + \frac{1}{10x^5} - \frac{1}{12x^6} + O(x^{-7})$$





# A motivating example

- The (central) **Delannoy polynomials**  $A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k$  satisfy  $A(-1) = 0, A(0) = 1$   
 $(n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}$

- Let  $B(n)$  be the 2nd solution with initial conditions  $B(0) = 0, B(1) = 1$ .

$$A(n) = 1, 1 + 2x, 1 + 6x + 6x^2, (1 + 2x)(1 + 10x + 10x^2), \dots$$

$$B(n) = 0, 1, \frac{3}{2}(1 + 2x), \frac{1}{6}(11 + 60x + 60x^2), \frac{5}{12}(1 + 2x)(5 + 42x + 42x^2), \dots$$

$$Q(n) := \frac{B(n)}{A(n)} = 0, \frac{1}{1+2x}, \frac{3(1+2x)}{2(1+6x+6x^2)}, \frac{11+60x+60x^2}{6(1+2x)(1+10x+10x^2)}, \dots$$

$$Q(n) - Q(n-1) = \frac{1}{1+2x}, \frac{1}{2(1+2x)(1+6x+6x^2)}, \dots \quad \frac{1}{nA(n)A(n-1)}$$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{2} \ln \left( 1 + \frac{1}{x} \right) = \sum_{n=1}^{\infty} \frac{1}{nA(n)A(n-1)}$$

$$Q(3) = \frac{1}{2x} - \frac{1}{4x^2} + \frac{1}{6x^3} - \frac{1}{8x^4} + \frac{1}{10x^5} - \frac{1}{12x^6} + O(x^{-7})$$



# Explore Apéry limits

- 1 Pick a binomial sum  $A(n)$ .  
Using creative telescoping, compute a recurrence satisfied by  $A(n)$ .

## Explore Apéry limits

- 1 Pick a binomial sum  $A(n)$ .  
Using creative telescoping, compute a recurrence satisfied by  $A(n)$ .
- 2 Compute the initial terms of a secondary solution  $B(n)$  to the recurrence.

## Explore Apéry limits

- 1 Pick a binomial sum  $A(n)$ .  
Using creative telescoping, compute a recurrence satisfied by  $A(n)$ .
- 2 Compute the initial terms of a secondary solution  $B(n)$  to the recurrence.
- 3 Try to identify  $\lim_{n \rightarrow \infty} B(n)/A(n)$ ,  
either numerically or as a power series in an additional parameter.

# Explore Apéry limits

- 1 Pick a binomial sum  $A(n)$ .  
Using creative telescoping, compute a recurrence satisfied by  $A(n)$ .
- 2 Compute the initial terms of a secondary solution  $B(n)$  to the recurrence.
- 3 Try to identify  $\lim_{n \rightarrow \infty} B(n)/A(n)$ ,  
either numerically or as a power series in an additional parameter.

EG  
HW

Use  $\sum_{k=0}^n \binom{n}{k} \binom{n-k}{k} x^k$  to rediscover the CF

$$\arctan(z) = \frac{z}{1+} \frac{1^2 z^2}{3+} \frac{2^2 z^2}{5+} \cdots \frac{n^2 z^2}{(2n+1)+} \cdots$$

# Explore Apéry limits

- 1 Pick a binomial sum  $A(n)$ .  
Using creative telescoping, compute a recurrence satisfied by  $A(n)$ .
- 2 Compute the initial terms of a secondary solution  $B(n)$  to the recurrence.
- 3 Try to identify  $\lim_{n \rightarrow \infty} B(n)/A(n)$ ,  
either numerically or as a power series in an additional parameter.

EG  
HW

Use  $\sum_{k=0}^n \binom{n}{k} \binom{n-k}{k} x^k$  to rediscover the CF

$$\arctan(z) = \frac{z}{1+} \frac{1^2 z^2}{3+} \frac{2^2 z^2}{5+} \cdots \frac{n^2 z^2}{(2n+1)+} \cdots$$

EG  
HW

Start with  $\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}^2 x^k$  and  $\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}^3 x^k$ .

Compare findings with those by Zudilin on simultaneous approximations to the logarithm, dilogarithm and trilogarithm.

# Explore Apéry limits

- 1 Pick a binomial sum  $A(n)$ .  
Using creative telescoping, compute a recurrence satisfied by  $A(n)$ .
- 2 Compute the initial terms of a secondary solution  $B(n)$  to the recurrence.
- 3 Try to identify  $\lim_{n \rightarrow \infty} B(n)/A(n)$ ,  
either numerically or as a power series in an additional parameter.

EG  
HW

Use  $\sum_{k=0}^n \binom{n}{k} \binom{n-k}{k} x^k$  to rediscover the CF

$$\arctan(z) = \frac{z}{1+} \frac{1^2 z^2}{3+} \frac{2^2 z^2}{5+} \cdots \frac{n^2 z^2}{(2n+1)+} \cdots$$

EG  
HW

Start with  $\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}^2 x^k$  and  $\sum_{k=0}^n \binom{n}{k} \binom{n+k}{k}^3 x^k$ .

Compare findings with those by Zudilin on simultaneous approximations to the logarithm, dilogarithm and trilogarithm.

EG  
bonus

For  $\sum_{k=0}^n \binom{n}{k}^2 \binom{3k}{n}$ , determine and prove the Apéry limits.

This is one of many cases conjectured by Almkvist, van Straten and Zudilin (2008) for CY DE's. Can we establish all these limits in a uniform fashion?

# Approaches to proving Apéry limits

Q

How to prove  $\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$ ?

1 Via explicit expressions:

(Apéry, '78)

$$B(n) = \frac{1}{6} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left( \sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right)$$



# Approaches to proving Apéry limits

Q

How to prove  $\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$ ?

1 Via explicit expressions:

(Apéry, '78)

$$B(n) = \frac{1}{6} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left( \sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right)$$

2 Via integral representations:

(Beukers, '79)

$$(-1)^n \int_0^1 \int_0^1 \int_0^1 \frac{x^n (1-x)^n y^n (1-y)^n z^n (1-z)^n}{(1 - (1-xy)z)^{n+1}} dx dy dz = A(n)\zeta(3) - 6B(n)$$

# Approaches to proving Apéry limits

Q

How to prove  $\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$ ?

1 Via explicit expressions:

(Apéry, '78)

$$B(n) = \frac{1}{6} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left( \sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right)$$

2 Via integral representations:

(Beukers, '79)

$$(-1)^n \int_0^1 \int_0^1 \int_0^1 \frac{x^n (1-x)^n y^n (1-y)^n z^n (1-z)^n}{(1 - (1-xy)z)^{n+1}} dx dy dz = A(n)\zeta(3) - 6B(n)$$

3 Via hypergeometric series representations:

(Gutnik, '79)

$$-\frac{1}{2} \sum_{t=1}^{\infty} R'_n(t) = A(n)\zeta(3) - 6B(n), \quad \text{where } R_n(t) = \left( \frac{(t-1) \cdots (t-n)}{t(t+1) \cdots (t+n)} \right)^2$$

# Approaches to proving Apéry limits

Q

How to prove  $\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$ ?

1 Via explicit expressions:

(Apéry, '78)

$$B(n) = \frac{1}{6} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left( \sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right)$$

2 Via integral representations:

(Beukers, '79)

$$(-1)^n \int_0^1 \int_0^1 \int_0^1 \frac{x^n (1-x)^n y^n (1-y)^n z^n (1-z)^n}{(1 - (1-xy)z)^{n+1}} dx dy dz = A(n)\zeta(3) - 6B(n)$$

3 Via hypergeometric series representations:

(Gutnik, '79)

$$-\frac{1}{2} \sum_{t=1}^{\infty} R'_n(t) = A(n)\zeta(3) - 6B(n), \quad \text{where } R_n(t) = \left( \frac{(t-1) \cdots (t-n)}{t(t+1) \cdots (t+n)} \right)^2$$

4 Via modular forms

(Beukers '87, Zagier '03, Yang '07)

# Approaches to proving Apéry limits

Q

How to prove  $\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{\zeta(3)}{6}$ ?

1 Via explicit expressions:

(Apéry, '78)

$$B(n) = \frac{1}{6} \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \left( \sum_{j=1}^n \frac{1}{j^3} + \sum_{m=1}^k \frac{(-1)^{m-1}}{2m^3 \binom{n}{m} \binom{n+m}{m}} \right)$$

2 Via integral representations:

(Beukers, '79)

$$(-1)^n \int_0^1 \int_0^1 \int_0^1 \frac{x^n (1-x)^n y^n (1-y)^n z^n (1-z)^n}{(1 - (1-xy)z)^{n+1}} dx dy dz = A(n)\zeta(3) - 6B(n)$$

3 Via hypergeometric series representations:

(Gutnik, '79)

$$-\frac{1}{2} \sum_{t=1}^{\infty} R'_n(t) = A(n)\zeta(3) - 6B(n), \quad \text{where } R_n(t) = \left( \frac{(t-1) \cdots (t-n)}{t(t+1) \cdots (t+n)} \right)^2$$

4 Via modular forms

(Beukers '87, Zagier '03, Yang '07)

5 Via continued fractions (for recurrences of order 2)

# Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 +} \frac{a_2}{b_2 +} \frac{a_3}{b_3 +} \dots := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

**THM** 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit:  $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

# Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

**THM** 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit:  $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

- Here,  $A(n), B(n)$  are the solutions to  $u_n = b_n u_{n-1} + a_n u_{n-2}$   
with  $A(-1) = 0, A(0) = 1$  and  $B(0) = 0, B(1) = a_1$ .

# Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 +} \frac{a_2}{b_2 +} \frac{a_3}{b_3 +} \dots := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

**THM** 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit:  $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

- Here,  $A(n), B(n)$  are the solutions to  $u_n = b_n u_{n-1} + a_n u_{n-2}$   
with  $A(-1) = 0, A(0) = 1$  and  $B(0) = 0, B(1) = a_1$ .

**proof** The  $n$ -th convergent is  $C_n := \frac{a_1}{b_1 +} \frac{a_2}{b_2 +} \dots \frac{a_n}{b_n} = \frac{B(n)}{A(n)}$ . □

# Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

**THM** 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit:  $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

- Here,  $A(n), B(n)$  are the solutions to  $u_n = b_n u_{n-1} + a_n u_{n-2}$   
with  $A(-1) = 0, A(0) = 1$  and  $B(0) = 0, B(1) = a_1$ .

**EG**

$$A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k \text{ solves } (n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}.$$

Let  $B(n)$  be the solution with  $B(0) = 0, B(1) = 1$ .



# Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} := \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

**THM** 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit:  $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

- Here,  $A(n), B(n)$  are the solutions to  $u_n = b_n u_{n-1} + a_n u_{n-2}$   
with  $A(-1) = 0, A(0) = 1$  and  $B(0) = 0, B(1) = a_1$ .

**EG**

$$A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k \text{ solves } (n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}.$$

Let  $B(n)$  be the solution with  $B(0) = 0, B(1) = 1$ .

$$\text{Hence, } n!A(n), n!B(n) \text{ solve } u_{n+1} = \frac{(2x+1)(2n+1)}{b_{n+1}} u_n - \frac{n^2}{a_{n+1}} u_{n-1}.$$

# Continued fractions and Apéry limits

$$C = \frac{a_1}{b_1 + \frac{a_2}{b_2 + \frac{a_3}{b_3 + \dots}}} \quad (b_n \neq 0)$$

**THM** 1-1 correspondence between CFs and order 2 recurrences, such that the value of the CF is an Apéry limit:  $C = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$

- Here,  $A(n), B(n)$  are the solutions to  $u_n = b_n u_{n-1} + a_n u_{n-2}$   
with  $A(-1) = 0, A(0) = 1$  and  $B(0) = 0, B(1) = a_1$ .

**EG**

$$A(n) = \sum_{k=0}^n \binom{n}{k} \binom{n+k}{k} x^k \text{ solves } (n+1)u_{n+1} = (2x+1)(2n+1)u_n - nu_{n-1}.$$

Let  $B(n)$  be the solution with  $B(0) = 0, B(1) = 1$ .

$$\text{Hence, } n!A(n), n!B(n) \text{ solve } u_{n+1} = \frac{(2x+1)(2n+1)}{b_{n+1}} u_n - \frac{n^2}{a_{n+1}} u_{n-1}.$$


Apéry limit and equivalent CF:

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \frac{1}{(2x+1) - \frac{1^2}{3(2x+1) - \frac{2^2}{5(2x+1) - \dots}}} = \frac{1}{2} \ln \left( 1 + \frac{1}{x} \right)$$

# Franel numbers

**DEF**  
Franel  
1894

$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$  are the (generalized) Franel numbers.




# Franel numbers

**DEF**  
Franel  
1894

$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$  are the (generalized) Franel numbers.

- $A^{(1)}(n) = 2^n$

$$u_{n+1} = 2u_n$$



# Franel numbers

DEF  
Franel  
1894


$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$  are the (generalized) Franel numbers.

- $A^{(1)}(n) = 2^n$

$$u_{n+1} = 2u_n$$

- $A^{(2)}(n) = \binom{2n}{n}$

$$(n+1)u_{n+1} = 2(2n+1)u_n$$



# Franel numbers

**DEF**  
Franel  
1894

$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$  are the (generalized) Franel numbers.

- $A^{(1)}(n) = 2^n$


$$u_{n+1} = 2u_n$$

- $A^{(2)}(n) = \binom{2n}{n}$

$$(n+1)u_{n+1} = 2(2n+1)u_n$$

- $A^{(3)}(n) = 1, 2, 10, 56, 346, 2252, 15184, 104960, 739162, \dots$

$$(n+1)^2 u_{n+1} = (7n^2 + 7n + 2)u_n + 8n^2 u_{n-1}$$



(Franel, 1894)

# Franel numbers

DEF  
Franel  
1894

$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$  are the (generalized) Franel numbers.

- $A^{(1)}(n) = 2^n$

$$u_{n+1} = 2u_n$$

- $A^{(2)}(n) = \binom{2n}{n}$

$$(n+1)u_{n+1} = 2(2n+1)u_n$$

- $A^{(3)}(n) = 1, 2, 10, 56, 346, 2252, 15184, 104960, 739162, \dots$


$$(n+1)^2 u_{n+1} = (7n^2 + 7n + 2)u_n + 8n^2 u_{n-1}$$

(Franel, 1894)

- $A^{(4)}(n) = 1, 2, 18, 164, 1810, 21252, 263844, 3395016, 44916498, \dots$

$$(n+1)^3 u_{n+1} = 2(2n+1)(3n^2 + 3n + 1)u_n + 4n(16n^2 - 1)u_{n-1}$$

(Franel, 1895)



# Franel numbers

**DEF**  
Franel  
1894

$A^{(s)}(n) = \sum_{k=0}^n \binom{n}{k}^s$  are the (generalized) Franel numbers.

- $A^{(1)}(n) = 2^n$

$$u_{n+1} = 2u_n$$

- $A^{(2)}(n) = \binom{2n}{n}$

$$(n+1)u_{n+1} = 2(2n+1)u_n$$

- $A^{(3)}(n) = 1, 2, 10, 56, 346, 2252, 15184, 104960, 739162, \dots$


$$(n+1)^2 u_{n+1} = (7n^2 + 7n + 2)u_n + 8n^2 u_{n-1}$$

(Franel, 1894)

- $A^{(4)}(n) = 1, 2, 18, 164, 1810, 21252, 263844, 3395016, 44916498, \dots$

$$(n+1)^3 u_{n+1} = 2(2n+1)(3n^2 + 3n + 1)u_n + 4n(16n^2 - 1)u_{n-1}$$

(Franel, 1895)



**CONJ**  
Franel,  
1895

The minimal recurrence for  $A^{(s)}(n)$  has order  $\lfloor \frac{s+1}{2} \rfloor$   
and degree  $s - 1$ .

(spoiler: the degree part is not true)



# Frenel's conjecture

**CONJ**  
Frenel,  
1895

The minimal recurrence for  $A^{(s)}(n)$  has order  $\lfloor \frac{s+1}{2} \rfloor$  and degree  $s - 1$ .

# Franel's conjecture

**CONJ**  
Franel,  
1895

The minimal recurrence for  $A^{(s)}(n)$  has order  $\lfloor \frac{s+1}{2} \rfloor$  and degree  $s - 1$ .

- Perlstadt '86: order 3 recurrences for  $s = 5, 6$  of degrees 6, 9 computed using MACSYMA and creative telescoping



# Franel's conjecture

**CONJ**  
Franel,  
1895

The minimal recurrence for  $A^{(s)}(n)$  has order  $\lfloor \frac{s+1}{2} \rfloor$  and degree  $s - 1$ .

- Perlstadt '86: order 3 recurrences for  $s = 5, 6$  of degrees 6, 9  
computed using MACSYMA and creative telescoping

**THM**  
Stoll '97

$A^{(s)}(n)$  satisfies a recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$ .

Cusick '89 also constructs such recurrences.



# Franel's conjecture

**CONJ**  
Franel,  
1895

The minimal recurrence for  $A^{(s)}(n)$  has order  $\lfloor \frac{s+1}{2} \rfloor$  and degree  $s - 1$ .

- Perlstadt '86: order 3 recurrences for  $s = 5, 6$  of degrees 6, 9 computed using MACSYMA and creative telescoping



**THM**  
Stoll '97

$A^{(s)}(n)$  satisfies a recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$ .

Cusick '89 also constructs such recurrences.



**CONJ**  
Bostan  
'21

The minimal recurrence for  $A^{(s)}(n)$  has order  $m = \lfloor \frac{s+1}{2} \rfloor$  and

$$\text{degree} = \begin{cases} \frac{1}{3}m(m^2 - 1) + 1, & \text{for even } s, \\ \frac{1}{3}m^3 - \frac{1}{2}m^2 + \frac{2}{3}m + \frac{(-1)^{m-1}}{4}, & \text{for odd } s. \end{cases}$$

If true, the degree grows like  $s^3/24$ .



# Franel's conjecture

**CONJ**

Franel,  
1895

The minimal recurrence for  $A^{(s)}(n)$  has order  $\lfloor \frac{s+1}{2} \rfloor$  and degree  $s - 1$ .

- Perlstadt '86: order 3 recurrences for  $s = 5, 6$  of degrees 6, 9 computed using MACSYMA and creative telescoping



**THM**

Stoll '97

$A^{(s)}(n)$  satisfies a recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$ .

Cusick '89 also constructs such recurrences.



**CONJ**

Bostan  
'21

The minimal recurrence for  $A^{(s)}(n)$  has order  $m = \lfloor \frac{s+1}{2} \rfloor$  and

$$\text{degree} = \begin{cases} \frac{1}{3}m(m^2 - 1) + 1, & \text{for even } s, \\ \frac{1}{3}m^3 - \frac{1}{2}m^2 + \frac{2}{3}m + \frac{(-1)^m - 1}{4}, & \text{for odd } s. \end{cases}$$

If true, the degree grows like  $s^3/24$ .



- Verified at least for  $s \leq 20$ .  
using MinimalRecurrence from the LREtools Maple package

# Franel's conjecture

**CONJ**  
Franel,  
1895

The minimal recurrence for  $A^{(s)}(n)$  has order  $\lfloor \frac{s+1}{2} \rfloor$  and degree  $s - 1$ .



- Perlstadt '86: order 3 recurrences for  $s = 5, 6$  of degrees 6, 9 computed using MACSYMA and creative telescoping

**THM**  
Stoll '97

$A^{(s)}(n)$  satisfies a recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$ .

Cusick '89 also constructs such recurrences.



**CONJ**  
Bostan  
'21

The minimal recurrence for  $A^{(s)}(n)$  has order  $m = \lfloor \frac{s+1}{2} \rfloor$  and

$$\text{degree} = \begin{cases} \frac{1}{3}m(m^2 - 1) + 1, & \text{for even } s, \\ \frac{1}{3}m^3 - \frac{1}{2}m^2 + \frac{2}{3}m + \frac{(-1)^{m-1}}{4}, & \text{for odd } s. \end{cases}$$

If true, the degree grows like  $s^3/24$ .



- Verified at least for  $s \leq 20$ .  
using MinimalRecurrence from the LREtools Maple package
- Goal: The minimal **telescoping** recurrence for  $A^{(s)}(n)$  has order  $\geq \lfloor \frac{s+1}{2} \rfloor$ .

# How to prove lower bounds for orders of recurrences?

EG

- $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$  : recurrence of order 2 (Apéry '78)

- $\sum_{k=0}^n \binom{n}{k}^s$  : recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$  (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

# How to prove lower bounds for orders of recurrences?

EG

- $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$  : recurrence of order 2 (Apéry '78)

- $\sum_{k=0}^n \binom{n}{k}^s$  : recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$  (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

- For fixed sequence, order 1 can be ruled out using Hyper, (Petkovšek '92)  
an algorithm to compute order 1 (right) factors of recurrence operators.



# How to prove lower bounds for orders of recurrences?

EG

- $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$  : recurrence of order 2 (Apéry '78)

- $\sum_{k=0}^n \binom{n}{k}^s$  : recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$  (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

- For fixed sequence, order 1 can be ruled out using Hyper, (Petkovšek '92)  
an algorithm to compute order 1 (right) factors of recurrence operators.
- For Franel numbers, order 1 can be ruled out for all  $s \geq 3$  (Yuan-Lu-Schmidt '08)  
using congruential properties.

# How to prove lower bounds for orders of recurrences?

EG

- $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$  : recurrence of order 2 (Apéry '78)

- $\sum_{k=0}^n \binom{n}{k}^s$  : recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$  (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

- For fixed sequence, order 1 can be ruled out using Hyper, (Petkovšek '92)  
an algorithm to compute order 1 (right) factors of recurrence operators.
- For Franel numbers, order 1 can be ruled out for all  $s \geq 3$  (Yuan-Lu-Schmidt '08)  
using congruential properties.
- There are algorithms for fixed recurrence operators (Beke 1894, Bronstein '94,  
for computing factors of differen(tial/ce) operators. Zhou-van Hoeij '19, ...)

# How to prove lower bounds for orders of recurrences?

EG

•  $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$  : recurrence of order 2 (Apéry '78)

•  $\sum_{k=0}^n \binom{n}{k}^s$  : recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$  (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

- For fixed sequence, order 1 can be ruled out using Hyper, (Petkovšek '92)  
an algorithm to compute order 1 (right) factors of recurrence operators.
- For Franel numbers, order 1 can be ruled out for all  $s \geq 3$  (Yuan-Lu-Schmidt '08)  
using congruential properties.
- There are algorithms for fixed recurrence operators (Beke 1894, Bronstein '94,  
for computing factors of differen(tial/ce) operators. Zhou-van Hoeij '19, ...)
- If  $A(n+1)/A(n) \rightarrow \mu$  for  $\mu \in \bar{\mathbb{Q}}$  of degree  $d$ , then  $A(n)$  cannot satisfy a  
recurrence over  $\mathbb{Q}$  of order less than  $d$ . (McIntosh '89)

# How to prove lower bounds for orders of recurrences?

EG

- $\sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2$ : recurrence of order 2 (Apéry '78)

- $\sum_{k=0}^n \binom{n}{k}^s$ : recurrence of order  $\lfloor \frac{s+1}{2} \rfloor$  (Stoll '97)

Could there be recurrences of lower order?

... and higher degree

- For fixed sequence, order 1 can be ruled out using Hyper, (Petkovšek '92)  
an algorithm to compute order 1 (right) factors of recurrence operators.
- For Franel numbers, order 1 can be ruled out for all  $s \geq 3$  (Yuan-Lu-Schmidt '08)  
using congruential properties.
- There are algorithms for fixed recurrence operators (Beke 1894, Bronstein '94,  
for computing factors of differen(tial/ce) operators. Zhou-van Hoeij '19, ...)
- If  $A(n+1)/A(n) \rightarrow \mu$  for  $\mu \in \bar{\mathbb{Q}}$  of degree  $d$ , then  $A(n)$  cannot satisfy a  
recurrence over  $\mathbb{Q}$  of order less than  $d$ . (McIntosh '89)

For Apéry numbers:  $\mu = (1 + \sqrt{2})^4$ .

For Franel numbers:  $\mu = 2^s$ . Not helpful!

# Solutions to the Franel number recurrences

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

# Solutions to the Franel number recurrences

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof  
outline

- Suppose:  $P(n, N) \binom{n}{k}^s = b(n, k+1) - b(n, k)$   
for a hypergeometric term  $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$ .



# Solutions to the Franel number recurrences

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof  
outline

- Suppose:  $P(n, N) \binom{n}{k-t}^s = b(n, k-t+1) - b(n, k-t)$   
for a hypergeometric term  $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$ .



# Solutions to the Franel number recurrences

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof  
outline

- 1 Suppose:  $P(n, N) \binom{n}{k-t}^s = b(n, k-t+1) - b(n, k-t)$   
for a hypergeometric term  $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$ .
- 2  $P(n, N) \sum_{k=\alpha}^{\beta-1} \binom{n}{k-t}^s = b(n, \beta-t) - b(n, \alpha-t)$   $b(n, t)$  entire for  $n \gg 0$





# Solutions to the Franel number recurrences

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof  
outline

- 1 Suppose:  $P(n, N) \binom{n}{k-t}^s = b(n, k-t+1) - b(n, k-t)$   
for a hypergeometric term  $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$ .
- 2  $P(n, N) \sum_{k=\alpha}^{\beta-1} \binom{n}{k-t}^s = b(n, \beta-t) - b(n, \alpha-t)$   $b(n, t)$  entire for  $n \gg 0$   
 $\alpha \ll 0$  and  $\beta \gg n$   $= O(t^s)$  since  $b(n, t) = \text{rat}(n, t) \binom{n}{t}^s$

□

# Solutions to the Franel number recurrences

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof  
outline

1 Suppose:  $P(n, N) \binom{n}{k-t}^s = b(n, k-t+1) - b(n, k-t)$   
for a hypergeometric term  $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$ .

2  $P(n, N) \sum_{k=\alpha}^{\beta-1} \binom{n}{k-t}^s = b(n, \beta-t) - b(n, \alpha-t)$   $b(n, t)$  entire for  $n \gg 0$   
 $\alpha \ll 0$  and  $\beta \gg n$   $= O(t^s)$  since  $b(n, t) = \text{rat}(n, t) \binom{n}{t}^s$

3  $P(n, N) \sum_{k=0}^n \binom{n}{k-t}^s = O(t^s)$  omitted terms are  $O(t^s)$

□

# Solutions to the Franel number recurrences

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

$$A^{(s)}(n, t) := \sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s} = \sum_{j \geq 0} A_j^{(s)}(n) t^{2j}$$

proof  
outline

① Suppose:  $P(n, N) \binom{n}{k-t}^s = b(n, k-t+1) - b(n, k-t)$   
for a hypergeometric term  $b(n, k) = \text{rat}(n, k) \binom{n}{k}^s$ .

②  $P(n, N) \sum_{k=\alpha}^{\beta-1} \binom{n}{k-t}^s = b(n, \beta-t) - b(n, \alpha-t)$   $b(n, t)$  entire for  $n \gg 0$   
 $\alpha \ll 0$  and  $\beta \gg n$   $= O(t^s)$  since  $b(n, t) = \text{rat}(n, t) \binom{n}{t}^s$

③  $P(n, N) \sum_{k=0}^n \binom{n}{k-t}^s = O(t^s)$  omitted terms are  $O(t^s)$

④  $A^{(s)}(n, t) = \left(\frac{\pi t}{\sin(\pi t)}\right)^s \sum_{k=0}^n \binom{n}{k-t}^s$  and so  $P(n, N)A^{(s)}(n, t) = O(t^s)$ .  $\square$

# Apéry limits for Franel numbers

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

# Apéry limits for Franel numbers

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

THM  
S-Zudilin  
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

# Apéry limits for Franel numbers

**THM**  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

**THM**  
S-Zudilin  
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

- To prove this, we show locally uniform convergence in  $t$  of

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s}}{\sum_{k=0}^n \binom{n}{k}^s} = \left( \frac{\pi t}{\sin(\pi t)} \right)^s.$$

# Apéry limits for Franel numbers

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

THM  
S-Zudilin  
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

- To prove this, we show locally uniform convergence in  $t$  of

$$\lim_{n \rightarrow \infty} \frac{\sum_{k=0}^n \binom{n}{k}^s \left[ \prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \right]^{-s}}{\sum_{k=0}^n \binom{n}{k}^s} = \left( \frac{\pi t}{\sin(\pi t)} \right)^s.$$

“poof” For large  $n$  and  $k \approx n/2$ ,

$$\prod_{j=1}^k \left(1 - \frac{t}{j}\right) \prod_{j=1}^{n-k} \left(1 + \frac{t}{j}\right) \approx \prod_{j=1}^{\infty} \left(1 - \frac{t}{j}\right) \left(1 + \frac{t}{j}\right) = \frac{\sin(\pi t)}{\pi t}.$$



# Apéry limits for Franel numbers

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

THM  
S-Zudilin  
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

$$\left( \frac{\pi t}{\sin(\pi t)} \right)^s = \left( \sum_{j=1}^{\infty} \left( 2 - \frac{1}{2^{2j-2}} \right) \zeta(2j) t^{2j} \right)^s = 1 + s\zeta(2) t^2 + \frac{s(5s+2)}{4} \zeta(4) t^4 + O(t^6)$$



# Apéry limits for Franel numbers

**THM**  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

**THM**  
S-Zudilin  
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

$$\left( \frac{\pi t}{\sin(\pi t)} \right)^s = \left( \sum_{j=1}^{\infty} \left( 2 - \frac{1}{2^{2j-2}} \right) \zeta(2j) t^{2j} \right)^s = 1 + s\zeta(2) t^2 + \frac{s(5s+2)}{4} \zeta(4) t^4 + O(t^6)$$

**EG**  
 $j=1$

$$\lim_{n \rightarrow \infty} \frac{B^{(s)}(n)}{A^{(s)}(n)} = \frac{1}{s(s+1)} s\zeta(2) = \frac{\zeta(2)}{s+1}$$

$$B^{(s)}(n) = \frac{A_1^{(s)}(n)}{A_1^{(s)}(1)} = 0, 1, \dots$$

$$A_1^{(s)}(1) = s(s+1)$$

# Apéry limits for Franel numbers

**THM**  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

**THM**  
S-Zudilin  
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

$$\left( \frac{\pi t}{\sin(\pi t)} \right)^s = \left( \sum_{j=1}^{\infty} \left( 2 - \frac{1}{2^{2j-2}} \right) \zeta(2j) t^{2j} \right)^s = 1 + s\zeta(2) t^2 + \frac{s(5s+2)}{4} \zeta(4) t^4 + O(t^6)$$

**EG**  
 $j=1$

$$\lim_{n \rightarrow \infty} \frac{B^{(s)}(n)}{A^{(s)}(n)} = \frac{1}{s(s+1)} s\zeta(2) = \frac{\zeta(2)}{s+1}$$

$$B^{(s)}(n) = \frac{A_1^{(s)}(n)}{A_1^{(s)}(1)} = 0, 1, \dots$$

- $s = 3, 4$  numerically observed by Cusick (1979)
- $s = 3$  proved by Zagier (2009)
- $s = 5$  conjectured by Almkvist, van Straten, Zudilin (2008)
- $s \geq 3$  conjectured by Chamberland-S (2020)

$$A_1^{(s)}(1) = s(s+1)$$

# Apéry limits for Franel numbers

**THM**  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

**THM**  
S-Zudilin  
'21

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

$$\left( \frac{\pi t}{\sin(\pi t)} \right)^s = \left( \sum_{j=1}^{\infty} \left( 2 - \frac{1}{2^{2j-2}} \right) \zeta(2j) t^{2j} \right)^s = 1 + s\zeta(2) t^2 + \frac{s(5s+2)}{4} \zeta(4) t^4 + O(t^6)$$

**EG**  
 $j = 1$

$$\lim_{n \rightarrow \infty} \frac{B^{(s)}(n)}{A^{(s)}(n)} = \frac{1}{s(s+1)} s\zeta(2) = \frac{\zeta(2)}{s+1}$$

$$B^{(s)}(n) = \frac{A_1^{(s)}(n)}{A_1^{(s)}(1)} = 0, 1, \dots$$

- $s = 3, 4$  numerically observed by Cusick (1979)
- $s = 3$  proved by Zagier (2009)
- $s = 5$  conjectured by Almkvist, van Straten, Zudilin (2008)
- $s \geq 3$  conjectured by Chamberland-S (2020)

$$A_1^{(s)}(1) = s(s+1)$$

**EG**  
 $j = 2$

$$\lim_{n \rightarrow \infty} \frac{C^{(s)}(n)}{A^{(s)}(n)} = \frac{12}{s(s+1)(s+2)(s+3)} \frac{s(5s+2)}{4} \zeta(4) \quad C^{(s)}(n) = \frac{A_2^{(s)}(n)}{A_2^{(s)}(1)} = 0, 1, \dots$$

- $s \geq 5$  conjectured by Chamberland-S (2020)

# Telescoping version of Franel's conjecture

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

# Telescoping version of Franel's conjecture

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for  $s \leq 30$ .

# Telescoping version of Franel's conjecture

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for  $s \leq 30$ .
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

# Telescoping version of Franel's conjecture

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for  $s \leq 30$ .
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by  $A_j^{(s)}(n) \in \mathbb{Q}$  if  $0 \leq 2j < s$ .

Here, and below, we assume that  $n$  is large enough.



# Telescoping version of Franel's conjecture

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for  $s \leq 30$ .
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by  $A_j^{(s)}(n) \in \mathbb{Q}$  if  $0 \leq 2j < s$ .

Here, and below, we assume that  $n$  is large enough.

2 The claim follows if these are linearly independent.





# Telescoping version of Franel's conjecture

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for  $s \leq 30$ .
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by  $A_j^{(s)}(n) \in \mathbb{Q}$  if  $0 \leq 2j < s$ .

Here, and below, we assume that  $n$  is large enough.

2 The claim follows if these are linearly independent.

3  $0 = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j A_j^{(s)}(n)$

$\lambda_j \in \mathbb{Q}$

□

# Telescoping version of Franel's conjecture

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for  $s \leq 30$ .
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

- 1 Any telescoping recurrence is solved by  $A_j^{(s)}(n) \in \mathbb{Q}$  if  $0 \leq 2j < s$ .

Here, and below, we assume that  $n$  is large enough.

- 2 The claim follows if these are linearly independent.

3  $0 = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j A_j^{(s)}(n) \implies 0 = \lim_{n \rightarrow \infty} \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j \frac{A_j^{(s)}(n)}{A^{(s)}(n)}$

$\lambda_j \in \mathbb{Q}$

□

# Telescoping version of Franel's conjecture

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for  $s \leq 30$ .
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by  $A_j^{(s)}(n) \in \mathbb{Q}$  if  $0 \leq 2j < s$ .

Here, and below, we assume that  $n$  is large enough.

2 The claim follows if these are linearly independent.

$$3 \quad 0 = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j A_j^{(s)}(n) \quad \implies \quad 0 = \lim_{n \rightarrow \infty} \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j \varphi_j \pi^{2j}$$

$\lambda_j \in \mathbb{Q}$    $\varphi_j \in \mathbb{Q}^\times$

□

# Telescoping version of Franel's conjecture

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- This implies Franel's conjecture on the exact order if the minimal-order recurrence is telescoping. True at least for  $s \leq 30$ .
- Order could be reduced by a different representation such as:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}$$

proof

1 Any telescoping recurrence is solved by  $A_j^{(s)}(n) \in \mathbb{Q}$  if  $0 \leq 2j < s$ .

Here, and below, we assume that  $n$  is large enough.

2 The claim follows if these are linearly independent.

$$3 \quad 0 = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j A_j^{(s)}(n) \quad \implies \quad 0 = \lim_{n \rightarrow \infty} \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = \sum_{j=0}^{\lfloor \frac{s-1}{2} \rfloor} \lambda_j \varphi_j \pi^{2j}$$

$\lambda_j \in \mathbb{Q}$    $\varphi_j \in \mathbb{Q}^\times$

4 Transcendence of  $\pi$  implies that all  $\lambda_j$  are zero. □

# When does creative telescoping fall short?

EG  
Paule,  
Schorn  
'95

$$\text{Consider } S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n}.$$



# When does creative telescoping fall short?

EG  
Paule,  
Schorn  
'95

Consider  $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n}$ .

- Any telescoping recurrence  $P(n, N)$  has order  $\geq d - 1$ :

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$



# When does creative telescoping fall short?

EG  
Paule,  
Schorn  
'95

Consider  $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$ .

- Any telescoping recurrence  $P(n, N)$  has order  $\geq d - 1$ :

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However,  $S_d(n) = (-d)^n$  satisfies  $NS_d(n) + dS_d(n) = 0$ .



# When does creative telescoping fall short?

EG  
Paule,  
Schorn  
'95

Consider  $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$ .

- Any telescoping recurrence  $P(n, N)$  has order  $\geq d - 1$ :

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However,  $S_d(n) = (-d)^n$  satisfies  $NS_d(n) + dS_d(n) = 0$ .



- Open problem: When does CT fall short?



# When does creative telescoping fall short?

EG  
Paule,  
Schorn  
'95

Consider  $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$ .

- Any telescoping recurrence  $P(n, N)$  has order  $\geq d - 1$ :

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However,  $S_d(n) = (-d)^n$  satisfies  $NS_d(n) + dS_d(n) = 0$ .



- Open problem: When does CT fall short?
- Can these cases be “fixed” by a different hypergeometric representation?

# When does creative telescoping fall short?

EG  
Paule,  
Schorn  
'95

Consider  $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$ .

- Any telescoping recurrence  $P(n, N)$  has order  $\geq d - 1$ :

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However,  $S_d(n) = (-d)^n$  satisfies  $NS_d(n) + dS_d(n) = 0$ .



- Open problem: When does CT fall short?
- Can these cases be “fixed” by a different hypergeometric representation?

EG  
Riese '01  
“creative  
symmetrizing”

Consider  $\sum_{k=1}^{2n} (-1)^k \binom{2n}{k}^2 \binom{2n}{k-1} = (-1)^n \frac{(3n)!}{n!^2(n-1)!(2n+1)}$ .



# When does creative telescoping fall short?

EG  
Paule,  
Schorn  
'95

Consider  $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$ .

- Any telescoping recurrence  $P(n, N)$  has order  $\geq d - 1$ :

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However,  $S_d(n) = (-d)^n$  satisfies  $NS_d(n) + dS_d(n) = 0$ .



- Open problem: When does CT fall short?
- Can these cases be “fixed” by a different hypergeometric representation?

EG  
Riese '01  
“creative  
symmetrizing”

Consider  $\sum_{k=1}^{2n} (-1)^k \binom{2n}{k}^2 \binom{2n}{k-1} = (-1)^n \frac{(3n)!}{n!^2(n-1)!(2n+1)}$ .

- CT produces order 2 recurrence on summand  $a(n, k)$ , but



# When does creative telescoping fall short?

EG  
Paule,  
Schorn  
'95

Consider  $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$ .

- Any telescoping recurrence  $P(n, N)$  has order  $\geq d - 1$ :

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However,  $S_d(n) = (-d)^n$  satisfies  $NS_d(n) + dS_d(n) = 0$ .



- Open problem: When does CT fall short?
- Can these cases be “fixed” by a different hypergeometric representation?

EG  
Riese '01  
“creative  
symmetrizing”

Consider  $\sum_{k=1}^{2n} (-1)^k \binom{2n}{k}^2 \binom{2n}{k-1} = (-1)^n \frac{(3n)!}{n!^2(n-1)!(2n+1)}$ .

- CT produces order 2 recurrence on summand  $a(n, k)$ , but
- order 1 on  $a(n, k) + a(n, 2n - k + 1) = \frac{2n - 2k + 1}{2n - k + 1} a(n, k)$ .



# When does creative telescoping fall short?

EG  
Paule,  
Schorn  
'95

Consider  $S_d(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} \binom{dk}{n} = (-d)^n$ .

- Any telescoping recurrence  $P(n, N)$  has order  $\geq d - 1$ :

$$P(n, N)(-1)^k \binom{n}{k} \binom{dk}{n} = b(n, k+1) - b(n, k)$$

- However,  $S_d(n) = (-d)^n$  satisfies  $NS_d(n) + dS_d(n) = 0$ .



- Open problem: When does CT fall short?
- Can these cases be “fixed” by a different hypergeometric representation?

EG  
Riese '01  
“creative  
symmetrizing”

Consider  $\sum_{k=1}^{2n} (-1)^k \binom{2n}{k}^2 \binom{2n}{k-1} = (-1)^n \frac{(3n)!}{n!^2(n-1)!(2n+1)}$ .

- CT produces order 2 recurrence on summand  $a(n, k)$ , but
- order 1 on  $a(n, k) + a(n, 2n - k + 1) = \frac{2n - 2k + 1}{2n - k + 1} a(n, k)$ .



“ Studying a huge number of practical applications one is tempted to conjecture that Zeilberger’s algorithm always returns the recurrence with minimal order. ”

Peter Paule, Markus Schorn, *Journal of Symbolic Computation*, 1995

# Modularity and Apéry limits

- Let  $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$  be Zagier's sporadic sequence **C**.

1, 3, 15, 93, ...

**THM**  
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left( \frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4)$$

$$x(\tau) = q - 4q^2 + 10q^3 + O(q^4)$$

$$q = e^{2\pi i \tau}$$

# Modularity and Apéry limits

- Let  $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$  be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

**THM**  
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left( \frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4)$$

$$x(\tau) = q - 4q^2 + 10q^3 + O(q^4)$$

$$q = e^{2\pi i \tau}$$

- Context:
  - $f(\tau)$  modular form of weight  $k$
  - $x(\tau)$  modular function
  - $y(x)$  such that  $y(x(\tau)) = f(\tau)$

Then  $y(x)$  satisfies a linear differential equation  $Ly = 0$  of order  $k + 1$ .

# Modularity and Apéry limits

- Let  $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$  be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

**THM**  
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left( \frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4)$$

$$x(\tau) = q - 4q^2 + 10q^3 + O(q^4)$$

$$q = e^{2\pi i \tau}$$

- Context:
  - $f(\tau)$  modular form of weight  $k$
  - $x(\tau)$  modular function
  - $y(x)$  such that  $y(x(\tau)) = f(\tau)$

Then  $y(x)$  satisfies a linear differential equation  $Ly = 0$  of order  $k + 1$ .

- Solutions to  $Ly = \text{rat}(x)$  are of the form  $y(x)$  times an Eichler integral of

$$h(\tau) = \left( \frac{Dx(\tau)}{x(\tau)} \right)^{k+1} \frac{\text{rat}(x(\tau))}{f(\tau)} \quad (\text{a modular form of weight } k+2) \quad (\text{Yang '07})$$

$$D = q \frac{d}{dq}$$

If  $\sum_{n \geq 1} c_n q^n$  is a modular form of weight  $k + 2$ , then  $\sum_{n \geq 1} \frac{c_n}{n^{k+1}} q^n$  is an Eichler integral.



# Modularity and Apéry limits

- Let  $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$  be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

**THM**  
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left( \frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4) \quad x(\tau) = q - 4q^2 + 10q^3 + O(q^4) \quad q = e^{2\pi i \tau}$$

- $F(x) := \sum_{n \geq 0} A(n)x^n \implies F(x(\tau)) = f(\tau)$
- $G(x) := \sum_{n \geq 0} B(n)x^n \implies G(x(\tau)) = f(\tau) \sum_{n \geq 1} \frac{\binom{-3}{n}}{n^2} \frac{q^n}{1+q^n}$

# Modularity and Apéry limits

- Let  $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$  be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

**THM**  
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left( \frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4) \quad x(\tau) = q - 4q^2 + 10q^3 + O(q^4) \quad q = e^{2\pi i \tau}$$

- $F(x) := \sum_{n \geq 0} A(n)x^n \implies F(x(\tau)) = f(\tau)$
- $G(x) := \sum_{n \geq 0} B(n)x^n \implies G(x(\tau)) = f(\tau) \sum_{n \geq 1} \frac{\binom{-3}{n}}{n^2} \frac{q^n}{1+q^n}$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \lim_{x \rightarrow \frac{1}{9}} \frac{G(x)}{F(x)}$$

characteristic roots 1, 9

$F(x), G(x)$  have radius of convergence  $R = \frac{1}{9}$ .

$G(x) - LF(x)$  has radius of convergence  $R = 1 > \frac{1}{9}$  for  $L = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$ .

# Modularity and Apéry limits

- Let  $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$  be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

**THM**  
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left( \frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4) \quad x(\tau) = q - 4q^2 + 10q^3 + O(q^4) \quad q = e^{2\pi i \tau}$$

- $F(x) := \sum_{n \geq 0} A(n)x^n \implies F(x(\tau)) = f(\tau)$
- $G(x) := \sum_{n \geq 0} B(n)x^n \implies G(x(\tau)) = f(\tau) \sum_{n \geq 1} \frac{\binom{-3}{n}}{n^2} \frac{q^n}{1+q^n}$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \lim_{x \rightarrow \frac{1}{9}} \frac{G(x)}{F(x)} = \lim_{\tau \rightarrow 0} \frac{G(x(\tau))}{F(x(\tau))}$$

characteristic roots 1, 9

$$x(\tau) = \frac{1}{9} \text{ for } \tau = 0 \text{ or } q = 1$$

$F(x), G(x)$  have radius of convergence  $R = \frac{1}{9}$ .

$G(x) - LF(x)$  has radius of convergence  $R = 1 > \frac{1}{9}$  for  $L = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$ .

# Modularity and Apéry limits

- Let  $A(n) = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}$  be Zagier's sporadic sequence **C**. 1, 3, 15, 93, ...

**THM**  
Zagier '09

$$\underbrace{\frac{\eta(2\tau)^6 \eta(3\tau)}{\eta(\tau)^3 \eta(6\tau)^2}}_{\text{modular form}} = \sum_{n \geq 0} A(n) \underbrace{\left( \frac{\eta(\tau)^4 \eta(6\tau)^8}{\eta(2\tau)^8 \eta(3\tau)^4} \right)^n}_{\text{modular function}}$$

$$f(\tau) = 1 + 3q + 3q^2 + 3q^3 + O(q^4) \quad x(\tau) = q - 4q^2 + 10q^3 + O(q^4) \quad q = e^{2\pi i \tau}$$

- $F(x) := \sum_{n \geq 0} A(n)x^n \implies F(x(\tau)) = f(\tau)$
- $G(x) := \sum_{n \geq 0} B(n)x^n \implies G(x(\tau)) = f(\tau) \sum_{n \geq 1} \frac{\binom{-3}{n}}{n^2} \frac{q^n}{1+q^n}$

$$\lim_{n \rightarrow \infty} \frac{B(n)}{A(n)} = \lim_{x \rightarrow \frac{1}{9}} \frac{G(x)}{F(x)} = \lim_{\tau \rightarrow 0} \frac{G(x(\tau))}{F(x(\tau))} = \lim_{q \rightarrow 1} \sum_{n \geq 1} \frac{\binom{-3}{n}}{n^2} \frac{q^n}{1+q^n} = \frac{1}{2} L_{-3}(2)$$

characteristic roots 1, 9

$$x(\tau) = \frac{1}{9} \text{ for } \tau = 0 \text{ or } q = 1$$

$F(x), G(x)$  have radius of convergence  $R = \frac{1}{9}$ .

$G(x) - LF(x)$  has radius of convergence  $R = 1 > \frac{1}{9}$  for  $L = \lim_{n \rightarrow \infty} \frac{B(n)}{A(n)}$ .

# Conclusions and some open questions

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover,  $A_j^{(s)}(n)$  with  $0 \leq 2j < s$  are linearly independent, so that any telescoping recurrence has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

# Conclusions and some open questions

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover,  $A_j^{(s)}(n)$  with  $0 \leq 2j < s$  are linearly independent, so that any telescoping recurrence has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- Cusick '89 and Stoll '97 construct recurrences for Franel numbers. Can these constructions produce telescoping recurrences?

# Conclusions and some open questions

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover,  $A_j^{(s)}(n)$  with  $0 \leq 2j < s$  are linearly independent, so that any telescoping recurrence has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- Cusick '89 and Stoll '97 construct recurrences for Franel numbers.  
Can these constructions produce telescoping recurrences?
- What can we learn from other families of binomial sums?  
Also, it would be nice to simplify some of the technical steps in the arguments.

# Conclusions and some open questions

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover,  $A_j^{(s)}(n)$  with  $0 \leq 2j < s$  are linearly independent, so that any telescoping recurrence has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- Cusick '89 and Stoll '97 construct recurrences for Franel numbers.  
Can these constructions produce telescoping recurrences?
- What can we learn from other families of binomial sums?  
Also, it would be nice to simplify some of the technical steps in the arguments.
- Can we (uniformly) establish the conjectural Apéry limits for CY DE's?



# Conclusions and some open questions

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover,  $A_j^{(s)}(n)$  with  $0 \leq 2j < s$  are linearly independent, so that any telescoping recurrence has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- Cusick '89 and Stoll '97 construct recurrences for Franel numbers.  
Can these constructions produce telescoping recurrences?
- What can we learn from other families of binomial sums?  
Also, it would be nice to simplify some of the technical steps in the arguments.
- Can we (uniformly) establish the conjectural Apéry limits for CY DE's?
- How to explicitly construct the solutions guaranteed by Perron's theorem?

# Conclusions and some open questions

THM  
S-Zudilin  
'21

Any telescoping recurrence for  $\sum_{k=0}^n \binom{n}{k}^s$  solved by  $A_j^{(s)}(n)$  if  $0 \leq 2j < s$ .  
(fine print: for large enough  $n$ )

The Apéry limits are:

$$\lim_{n \rightarrow \infty} \frac{A_j^{(s)}(n)}{A^{(s)}(n)} = [t^{2j}] \left( \frac{\pi t}{\sin(\pi t)} \right)^s \in \pi^{2j} \mathbb{Q}_{>0}$$

Moreover,  $A_j^{(s)}(n)$  with  $0 \leq 2j < s$  are linearly independent, so that any telescoping recurrence has order at least  $\lfloor \frac{s+1}{2} \rfloor$ .

- Cusick '89 and Stoll '97 construct recurrences for Franel numbers.  
Can these constructions produce telescoping recurrences?
- What can we learn from other families of binomial sums?  
Also, it would be nice to simplify some of the technical steps in the arguments.
- Can we (uniformly) establish the conjectural Apéry limits for CY DE's?
- How to explicitly construct the solutions guaranteed by Perron's theorem?
- Can we explain when CT falls short? And algorithmically "fix" this issue?

# THANK YOU!

Slides for this talk will be available from my website:

<http://arminstraub.com/talks>



**M. Chamberland, A. Straub**

*Apéry limits: Experiments and proofs*

American Mathematical Monthly, Vol. 128, Nr. 9, 2021, p. 811-824



**A. Straub, W. Zudilin**

*Sums of powers of binomials, their Apéry limits, and Franel's suspicions*

International Mathematics Research Notices, to appear, 2022. arXiv:2112.09576