

**Example 53. (review)** Consider the sequence  $a_n$  defined by  $a_{n+2} = a_{n+1} + 2a_n$  and  $a_0 = 1$ ,  $a_1 = 8$ .

- Determine the first few terms of the sequence.
- Find a formula for  $a_n$ .
- Determine  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}$ .

**Solution.**

- $a_2 = 10$ ,  $a_3 = 26$
- The recursion can be written as  $p(N)a_n = 0$  where  $p(N) = N^2 - N - 2$  has roots  $2, -1$ .  
Hence,  $a_n = C_1 2^n + C_2 (-1)^n$  and we only need to figure out the two unknowns  $C_1, C_2$ . We can do that using the two initial conditions:  $a_0 = C_1 + C_2 = 1$ ,  $a_1 = 2C_1 - C_2 = 8$ .  
Solving, we find  $C_1 = 3$  and  $C_2 = -2$  so that, in conclusion,  $a_n = 3 \cdot 2^n - 2(-1)^n$ .
- It follows from the formula  $a_n = 3 \cdot 2^n - 2(-1)^n$  that  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = 2$ .

**Comment.** In fact, this already follows from  $a_n = C_1 2^n + C_2 (-1)^n$  provided that  $C_1 \neq 0$ . Since  $a_n = C_2 (-1)^n$  (the case  $C_1 = 0$ ) is not compatible with  $a_0 = 1$ ,  $a_1 = 8$ , we can conclude  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = 2$  without computing the actual values of  $C_1$  and  $C_2$ .

## Crash course: Eigenvalues and eigenvectors

If  $Ax = \lambda x$  (and  $x \neq 0$ ), then  $x$  is an **eigenvector** of  $A$  with **eigenvalue**  $\lambda$  (just a number).

Note that, for the equation  $Ax = \lambda x$  to make sense,  $A$  needs to be a square matrix (i.e.  $n \times n$ ).

Key observation:

$$\begin{aligned} Ax &= \lambda x \\ \iff Ax - \lambda x &= 0 \\ \iff (A - \lambda I)x &= 0 \end{aligned}$$

This homogeneous system has a nontrivial solution  $x$  if and only if  $\det(A - \lambda I) = 0$ .

To find eigenvectors and eigenvalues of  $A$ :

- First, find the eigenvalues  $\lambda$  by solving  $\det(A - \lambda I) = 0$ .

$\det(A - \lambda I)$  is a polynomial in  $\lambda$ , called the **characteristic polynomial** of  $A$ .

- Then, for each eigenvalue  $\lambda$ , find corresponding eigenvectors by solving  $(A - \lambda I)x = 0$ .

**Example 54.** Determine the eigenvalues and eigenvectors of  $A = \begin{bmatrix} 8 & -10 \\ 5 & -7 \end{bmatrix}$ .

**Solution.** The characteristic polynomial is:

$$\det(A - \lambda I) = \det\left(\begin{bmatrix} 8-\lambda & -10 \\ 5 & -7-\lambda \end{bmatrix}\right) = (8-\lambda)(-7-\lambda) + 50 = \lambda^2 - \lambda - 6 = (\lambda-3)(\lambda+2)$$

Hence, the eigenvalues are  $\lambda = 3$  and  $\lambda = -2$ .

- To find an eigenvector for  $\lambda = 3$ , we need to solve  $\begin{bmatrix} 5 & -10 \\ 5 & -10 \end{bmatrix} \mathbf{x} = \mathbf{0}$ .  
Hence,  $\mathbf{x} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$  is an eigenvector for  $\lambda = 3$ .
- To find an eigenvector for  $\lambda = -2$ , we need to solve  $\begin{bmatrix} 10 & -10 \\ 5 & -5 \end{bmatrix} \mathbf{x} = \mathbf{0}$ .  
Hence,  $\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  is an eigenvector for  $\lambda = -2$ .

**Check!**  $\begin{bmatrix} 8 & -10 \\ 5 & -7 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 6 \\ 3 \end{bmatrix} = 3 \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix}$  and  $\begin{bmatrix} 8 & -10 \\ 5 & -7 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \end{bmatrix} = -2 \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

On the other hand, a random other vector like  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$  is not an eigenvector:  $\begin{bmatrix} 8 & -10 \\ 5 & -7 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \begin{bmatrix} -12 \\ -9 \end{bmatrix} \neq \lambda \begin{bmatrix} 1 \\ 2 \end{bmatrix}$ .

**Example 55. (homework)** Determine the eigenvalues and eigenvectors of  $A = \begin{bmatrix} 1 & -6 \\ 1 & -4 \end{bmatrix}$ .

**Solution. (final answer only)**  $\mathbf{x} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$  is an eigenvector for  $\lambda = -2$ , and  $\mathbf{x} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$  is an eigenvector for  $\lambda = -1$ .